Reduction of Compound Lotteries with

Objective Probabilities: Theory and Evidence

by

Glenn W. Harrison, Jimmy Martínez-Correa and J. Todd Swarthout †

March 2012

ABSTRACT.

The reduction of compound lotteries (ROCL) has assumed a central role in the evaluation of behavior towards risk and uncertainty. We present experimental evidence on its validity in the domain of objective probabilities. Our experiment explicitly recognizes the impact that the random lottery incentive mechanism payment procedure may have on preferences, and so we collect data using both "1-in-1" and "1-in-K" payment procedures, where K>1. We do not find violations of ROCL when subjects are presented with only one choice that is played for money. However, when individuals are presented with many choices and random lottery incentive mechanism is used to select one choice for payoff, we do find violations of ROCL. These results are supported by both non-parametric analysis of choice patterns, as well as structural estimation of latent preferences. We find evidence that the model that best describes behavior when subjects make only one choice is the Rank-Dependent Utility model. When subjects face many choices, their behavior is better characterized by our source-dependent version of the Rank-Dependent Utility model which can account for violations of ROCL. We conclude that payment protocols can create distortions in experimental tests of basic axioms of decision theory.

[†] Department of Risk Management & Insurance and Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, USA (Harrison); Department of Risk Management & Insurance, Robinson College of Business, Georgia State University, USA (Martínez-Correa); and Department of Economics, Andrew Young School of Policy Studies, Georgia State University, USA (Swarthout). E-mail contacts: gharrison@gsu.edu, jimmigan@gmail.com and swarthout@gsu.edu.

Table of Contents

1. Theory
B. Experimental Payment Protocols -5-
2. Experiment
3. Non-Parametric Analysis of Choice Patterns11-
A. Choice Patterns Where ROCL Predicts Indifference11- B. Choice Patterns Where ROCL Predicts Consistent Choices14-
4. Estimated Preferences from Observed Choices -17- A. Econometric Specification -17- B. Estimates -23-
5. Conclusions
References -40-
Appendix A: Instructions (NOT FOR PUBLICATION)
Appendix B: Parameters
Appendix C: Related Literature (NOT FOR PUBLICATION)62-
Appendix D: Nonparametric Tests (NOT FOR PUBLICATION)65-
Appendix E: Additional Econometric Analysis (NOT FOR PUBLICATION)74-
Appendix F: Detailed Binomial Test Results (NOT FOR PUBLICATION)80-
Appendix G: Detailed Fisher Exact Test Results (NOT FOR PUBLICATION)82-
Appendix H: Detailed McNemar Test Results (NOT FOR PUBLICATION)85-
Appendix I: Detailed Wald Test Results for Predictions (NOT FOR PUBLICATION)93-

The reduction of compound lotteries has assumed a central role in the evaluation of behavior towards risk, uncertainty and ambiguity. We present experimental evidence on its validity in domains defined over objective probabilities, as a prelude to evaluating it over subjective probabilities.

Because of the attention paid to violations of the Independence Axiom, it is noteworthy that early formal concerns with the possibility of a "utility or disutility for gambling" centered around the Reduction of Compound Lotteries (ROCL) axiom.¹ Von Neumann and Morgenstern [1953, p. 28] commented on the possibility of allowing for a (dis)utility of gambling component in their preference representation:

Do not our postulates introduce, in some oblique way, the hypotheses which bring in the mathematical expectation [of utility]? More specifically: May there not exist in an individual a (positive or negative) utility of the mere act of 'taking a chance,' of gambling, which the use of the mathematical expectation obliterates? How did our axioms (3:A)-(3:C) get around this possibility? As far as we can see, our postulates (3:A)-(3:C) do not attempt to avoid it. Even the one that gets closest to excluding the 'utility of gambling' - (3:C:b)- seems to be plausible and legitimate - unless a much more refined system of psychology is used than the one now available for the purposes of economics [...] Since (3:A)-(3:C) secure that the necessary construction [of utility] can be carried out, concepts like a 'specific utility of gambling' cannot be formulated free of contradiction on this level.

On the very last page of their *magnus opus*, von Neumann and Morgenstern [1953; p. 632] propose that if their postulate (3:C:b), which is the ROCL, is relaxed, one could indeed allow for a specific utility for the act of gambling:

It seems probable, that the really critical group of axioms is (3:C) - or, more specifically, the axiom (3:C:b). This axiom expresses the combination rule for multiple chance alternatives, and it is plausible, that a specific utility or disutility of gambling can only exist if this simple combination rule is abandoned. Some change of the system [of

¹ The issue of the (dis)utility of gambling goes back at least as far as Pascal, who argued in his *Pensées* that "people distinguish between the pleasure or displeasure of chance (uncertainty) and the objective evaluation of the worth of the gamble from the perspective of its consequences" (see Luce and Marley [2000; p. 102]). Referring to the ability of bets to elicit beliefs, Ramsey [1926] claims that "[t]his method I regard as fundamentally sound; but it suffers from being insufficiently general, and from being necessarily inexact. It is inexact partly [...] because the person may have a special eagerness or reluctance to bet, because he either enjoys or dislikes excitement or for any other reason, e.g. to make a book. The difficulty is like that of separating two different cooperating forces" (from the reprint in Kyburg and Smokler [1964; p. 73]).

axioms] (3:A)-(3:B), at any rate involving the abandonment or at least a radical modification of (3:C:b), may perhaps lead to a mathematically complete and satisfactory calculus of utilities which allows for the possibility of a specific utility or disutility of gambling. It is hoped that a way will be found to achieve this, but the mathematical difficulties seem to be considerable.

Thus, the relaxation of the ROCL axiom opens the door to the possibility of having a distinct (dis)utility for the act of gambling with objective probabilities.² Fellner [1961][1963] and Smith [1969] used similar reasoning to offer an explanation for several of the Ellsberg [1961] paradoxes.

This argument rests on the hypothesis that subjects potentially view simple and compound random processes differently. If this hypothesis is true, it could explain why people prefer risky over ambiguous gambles in the thought experiments of Ellsberg [1961]. Fellner [1961][1963] and Smith [1969] believed that if a subject could exhibit utility or disutility of gambling she may also use different utility functions to make decisions under different processes. Smith [1969] went further and explicitly conjectured that a compound lottery defined over *objective* probabilities, and its actuarially-equivalent lottery over *objective* probabilities, might be viewed by decision makers as two different random processes. In fact, he proposed a preference representation that allowed people to have different utility functions for different random processes. We use this conjectured preference representation to test for violations of ROCL.

One fundamental methodological problem with tests of the ROCL assumption, whether or not the context is objective or subjective probabilities, is that one cannot use incentives for decision makers that rely on the validity of ROCL. This means, in effect, that experiments must be conducted in which a subject has one, and only one, choice.³ Apart from the expense and time of collecting data at such a

² Of course, it is of some comfort to the egos of modern theorists that no less than von Neumann and Morganstern at least viewed it as a serious mathematical challenge.

³ One alternative is to present the decision maker with several tasks at once and evaluate the portfolio chosen, or to present the decision maker with several tasks in sequence and account for wealth effects. Neither is attractive, since they each raise a number of (fascinating) theoretical confounds to the interpretation of observed behavior. One uninteresting alternative is not to pay the decision maker for the outcomes of the task.

pace, this also means that all evaluations have to be on a between-subjects basis, implying the necessity of modeling assumptions about heterogeneity in behavior.

In sections 1 and 2 we define the theory and experimental tasks used to examine ROCL in the context of objective probabilities. In section 3 and 4 we present evidence from our experiment. We find *no violations of ROCL* when subjects are presented with one and only one choice, and that their behavior is better characterized by the Rank-Dependent Utility model (RDU) rather than Expected Utility Theory (EUT). However, we do find *violations of ROCL* when many choices are given to each subject and the random lottery incentive mechanism (RLIM) is used as the payment protocol. Under RLIM, behavior is better characterized by our source-dependent version of RDU that can account for violations of ROCL. Section 5 draws conclusions for modeling, experimental design, and inference about decision making.

1. Theory

A. Basic Axioms

Following Segal [1988][1990][1992], we distinguish between three axioms. In words, the **Reduction of Compound Lotteries** axiom states that a decision-maker is indifferent between a compound lottery and the actuarially-equivalent simple lottery in which the probabilities of the two stages of the compound lottery have been multiplied out. To use the language of Samuelson [1952; p.671], the former generates a *compound income-probability-situation*, and the latter defines an *associated income-probability-situation*, and that "...only algebra, not human behavior, is involved in this definition."

To state this more explicitly, with notation to be used to state all axioms, let X, Y and Z denote simple lotteries, A and B denote compound lotteries, > express strict preference, and ~ express indifference. Then the ROCL axiom says that A ~ X if the probabilities and prizes in X are the actuarially-equivalent probabilities and prizes from A. Thus if A is the compound lottery that pays "double or nothing" from the outcome of the lottery that pays \$10 if a coin flip is a head and \$2 if the

coin flip is a tail, then X would be the lottery that pays \$20 with probability $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$, \$4 with probability $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$, and nothing with probability $\frac{1}{2}$. From an observational perspective, one must see choices between compound lotteries and actuarially-equivalent simple lotteries to test ROCL.

The **Compound Independence Axiom** (CIA) states that two compound lotteries, each formed from a simple lottery by adding a positive common lottery with the same probability, will exhibit the same preference ordering as the simple lotteries. This is a statement that the ordering of the two constructed compound lotteries will be the same as the ordering of the different simple lotteries that distinguish the compound lotteries, provided that the common prize in the compound lotteries is the same and has the same (compound lottery) probability. It says nothing about how the compound lotteries are to be evaluated, and in particular *it does not assume ROCL*. It only restricts the preference *ordering* of the two constructed compound lotteries to match the preference *ordering* of the original simple lotteries.

The CIA says that if A is the compound lottery giving the simple lottery X with probability α and the simple lottery Z with probability (1- α), and B is the compound lottery giving the simple lottery Y with probability α and the simple lottery Z with probability (1- α), then A > B iff $X > Y \forall \alpha \in (0,1)$. The construction of the two compound lotteries A and B has the "independence axiom" cadence of the common prize Z with a common probability (1- α), but the implication is only that the ordering of the compound and constituent simple lotteries are the same.⁴

Finally, the **Mixture Independence Axiom** (MIA) says that the preference ordering of two simple lotteries must be the same as the actuarially-equivalent simple lottery formed by adding a

⁴ For example, Segal [1992; p.170] defines the CIA by assuming that the second-stage lotteries are replaced by their certainty-equivalent, "throwing away" information about the second-stage probabilities before one examines the first-stage probabilities at all. Hence one cannot then *define* the actuarially-equivalent simple lottery, by construction, since the informational bridge to that calculation has been burnt. The certainty-equivalent could have been generated by any model of decision making under risk, such as RDU or Prospect Theory.

common outcome in a compound lottery of each of the simple lotteries, where the common outcome has the same value and the same (compound lottery) probability. So stated, it is clear that the MIA strengthens the CIA by making a definite statement that the constructed compound lotteries are to be evaluated in a way that is ROCL-consistent. Construction of the compound lottery in the MIA is actually implicit: the axiom only makes observable statements about two pairs of simple lotteries. To restate Samuelson's point about the definition of ROCL, the experimenter testing the MIA could have constructed the associated income-probability-situation without knowing the risk preferences of the individual (although the experimenter would need to know how to multiply).

The MIA says that X > Y iff the actuarially-equivalent simple lottery of $\alpha X + (1-\alpha)Z$ is strictly preferred to the actuarially-equivalent simple lottery of $\alpha Y + (1-\alpha)Z$, $\forall \alpha \in (0,1)$. The verbose language used to state the axiom makes it clear that MIA embeds ROCL into the usual independence axiom construction with a common prize Z and a common probability $(1-\alpha)$ for that prize.

The reason these three axioms are important is that the failure of MIA does not imply the failure of CIA and ROCL. It does imply the failure of one or the other, but it is far from obvious which one. Indeed, one could imagine some individuals or task domains where only CIA might fail, only ROCL might fail, or both might fail. Because specific types of failures of ROCL lie at the heart of many important models of decision-making under uncertainty and ambiguity, it is critical to keep the axioms distinct as a theoretical *and* experimental matter.

B. Experimental Payment Protocols

Turning now to experimental procedures, as a matter of theory the most popular payment protocol assumes the validity of MIA. This payment protocol is called the **Random Lottery Incentive Mechanism** (RLIM). It entails the subject undertaking K>1 tasks and then one of the K choices being selected at random to be played out. Typically, and without loss of generality, assume that the selection

of the k^{th} task to be played out uses a uniform distribution over the K tasks. Since the other K-1 tasks will generate a payoff of zero, the payment protocol can be seen as a compound lottery that assigns probability $\alpha = 1/K$ to the selected task and $(1-\alpha) = (1-(1/K))$ to the other K-1 tasks as a whole. If the task consists of binary choices between simple lotteries X and Y, then the RLIM can be immediately seen to entail an application of MIA, where Z = U(\$0) and $(1-\alpha) = (1-(1/K))$, for the utility function $U(\cdot)$. Hence, under MIA, the preference ordering of X and Y is independent of all of the choices in the other tasks (Holt [1986]).

The need to assume the MIA can be avoided by setting K=1, and asking each subject to answer one binary choice task for payment. Unfortunately, this comes at the cost of another assumption if one wants to compare choice patterns over two simple lottery pairs, as in most of the popular tests of EUT such as the Allais Paradox and Common Ratio test: the assumption that risk preferences across subjects are the same. This is a strong assumption, obviously, and one that leads to inferential tradeoffs in terms of the "power" of tests of EUT relying on randomization that will vary with sample size. Sadly, plausible estimates of the degree of heterogeneity in the typical population imply massive sample sizes for reasonable power, well beyond those of most experiments.

The assumption of homogeneous preferences can be diluted, however, by changing it to a conditional form: that risk preferences are homogeneous conditional on a finite set of observable characteristics.⁵ Although this sounds like an econometric assumption, and it certainly has statistical implications, it is as much a matter of (operationally meaningful) theory as formal statements of the CIA, ROCL and MIA.

⁵ Another way of diluting the assumption is to posit some (flexible) parametric form for the distribution of risk attitudes in the population, and use econometric methods that allow one to estimate the extent of that unobserved heterogeneity across individuals. Tools for this "random coefficients" approach to estimating non-linear preference functionals are developed in Andersen, Harrison, Hole, Lau and Rutström [2010].

2. Experiment

A. Lottery Parameters

We designed our battery of lotteries to allow for specific types of comparisons needed for testing ROCL. Beginning with a given *simple* (S) lottery and *compound* (C) lottery, we next create an *actuarially-equivalent* (AE) lottery from the C lottery, and then we construct three pairs of lotteries: a S-C pair, a S-AE pair, and an AE-C pair. By repeating this process 15 times, we create a battery of lotteries consisting of 15 S-C pairs shown in Table B2, 15 S-AE pairs shown in Table B3, and 10 AE-C pairs⁶ shown in Table B4. See Appendix B for additional information regarding the creation of these lotteries.

Figure 1 displays the coverage of lotteries in the Marschak-Machina triangle, covering all of the contexts used.⁷ Probabilities were drawn from 0, ½, ½, ¾ and 1, and the final prizes from \$0, \$10, \$20, \$35 and \$70. We use the familiar "Double or Nothing" (DON) procedure for creating compound lotteries. So, the first-stage prizes displayed in a compound lottery were drawn from \$5, \$10, \$17.50 and \$35, and then the second-stage DON procedure yields the set of final prizes given above.

The majority of our compound lotteries use a conditional version of DON in the sense that the initial lottery will trigger the double or nothing option that the subject will face only if a particular outcome is realized in the initial lottery. For example, consider the compound lottery formed by an initial lottery that pays \$10 and \$20 with equal probability and the option of playing DON if the outcome of the initial lottery is \$10, implying a payoff of \$20 or \$0 with equal chance *if the DON stage is reached.* If the initial outcome is \$20, there is no DON option beyond that. The right panel of Figure 2 shows a tree representation of the latter compound lottery where the initial lottery is depicted in the first stage and the DON lottery is depicted in the second stage of the compound lottery if reached. The left

⁶ The lottery battery contains only 10 AE-C lottery pairs because some of the 15 S-C lottery pairs shared the same compound lottery.

⁷ Decision screens were presented to subjects in color. Black borders were added to each pie slice in Figures 1, 2 and 3 to facilitate black-and-white viewing.

panel of Figure 2 shows the corresponding actuarially-equivalent simple lottery which offers \$20 with probability 3/4 and \$0 with probability 1/4.

The conditional DON lottery allows us to obtain good coverage in terms of prizes and probabilities and to maintain a simple random processes for the initial lottery and the DON option. One can construct a myriad of compound lotteries with only two components: (1) initial lotteries that pay two outcomes with 50:50 odds or pay a given stake with certainty; and (2) a conditional DON which pays double a predetermined amount with 50% probability or nothing with equal chance. Using only the unconditional DON option would impose an *a priori* restriction on the coverage within the Marschak-Machina triangle.

B. Experimental Procedures

We implement two between-subjects treatments. We call one treatment "Pay 1-in-1" (1-in-1) and the other "Pay 1-in-40" (1-in-40). Table 1 summarizes our experimental design and the sample size of subjects and choices in each treatment.

In the 1-in-1 treatment, each subject faces a single choice over two lotteries. The lottery pair presented to each subject is randomly selected from the battery of 40 lottery pairs. The lottery chosen by the subject is then played out and the subject receives the realized monetary outcome. There are no other salient tasks, before or after a subject's binary choice, that affect the outcome. Further, there is no other activity that may contribute to learning about decision making in this context.

In the 1-in-40 treatment, each subject faces choices over all 40 lottery pairs, with the order of the pairs randomly shuffled for each subject. After all choices have been made, one choice is randomly selected for payment using the RLIM, with each choice having a 1-in-40 chance of being selected. The selected choice is then played out and the subject receives the realized monetary outcome, again with no other salient tasks. This treatment is potentially different from the 1-in-1 treatment in the absence of

ROCL, since the RLIM induces a compound lottery consisting of a 1-in-40 chance for each of the 40 chosen lotteries to be selected for payment.

The general procedures during an experiment session were as follows. Upon arrival at the laboratory, each subject drew a number from a box which determined random seating position within the laboratory. After being seated and signing the informed consent document, subjects were given printed instructions and allowed sufficient time to read these instructions. Once subjects had finished reading the instructions, an experimenter at the front of the room read aloud the instructions, word for word. Then the randomizing devices were explained and projected onto the front screen and three large flat-screen monitors spread throughout the laboratory. The subjects were then presented with lottery choices, followed by a non-salient demographic questionnaire that did not affect final payoffs. Next, each subject was approached by an experimenter who would provide dice so for the subject to roll and determine her own payoff. If a DON stage was reached, a subject would flip a U.S. quarter dollar coin to determine the final outcome of the lottery. Finally, subjects then left the laboratory and were privately paid their earnings: a \$7.50 participation payment in addition to the monetary outcome of the realized lottery.

We used software created in Visual Basic .NET to present lotteries to subjects and record their choices. Figure 3 shows an example of the subject display of an AE-C lottery pair. The first and second stages of the compound lottery, like the one depicted in Figure 2, are presented as an initial lottery, represented by the pie on the right of Figure 3, that has a DON option identified by text. The pie chart on the left of Figure 3 shows the AE lottery of the paired C lottery on the right. Figure 4 shows an example of the subject display of a S-C lottery pair, and Figure 5 shows an example of the subject

⁸ Appendix A provides complete subject instructions.

⁹ Only physical randomizing devices were used, and these devices were demonstrated prior to any decisions. In the 1-in-40 treatment, two 10-sided dice were rolled by each subject until a number between 1 and 40 came up to select the relevant choice for payment. Subjects in both treatments would roll the two 10-sided dice (a second roll in the case of the 1-in-40 treatment) to determine the outcome of the chosen lottery.

display of a S-AE lottery pair.

C. Evaluation of Hypotheses

If the subjects in both treatments have the same risk preferences and behavior is consistent with ROCL, we should see the same pattern of decisions for comparable lottery pairs across the two treatments. The same pattern should also be observed as one characterizes heterogeneity of individual preferences towards risk, although these inferences depend on the validity of the manner in which heterogeneity is modeled.

Nothing here assumes that behavior is characterized by EUT. The validity of EUT requires both ROCL and CIA, and the validity of ROCL does not imply the validity of CIA. So when we say that risk preferences should be the same in the two treatments under ROCL, these are simply statements about the Arrow-Pratt risk premium, and not about how that is decomposed into explanations that rely on diminishing marginal utility or probability weighting. We later analyze the decomposition of the risk premium as well as the nature of any violation of ROCL.

Our method of evaluation is twofold. First, we use non-parametric tests to evaluate the choice patterns of subjects. Our experimental design allows us to evaluate ROCL using choice patterns in two ways: (1) directly examine choice patterns in AE-C lottery pairs where ROCL predicts indifference; and (2) examine the choice patterns across the linked S-C and S-AE lottery pairs. We have 15 tests, one for each linked pair of lottery pairs, as well as a pooled test over all 15 pairs of pairs. We are agnostic as to the choice pattern itself: if subjects have a clear preference for S over C in a given lottery pair, then under ROCL we should see the same preference for the identical S over the AE in the linked lottery pair.

For our second method of evaluation of ROCL, we estimate structural models of risk preferences and test if the risk preference parameters depend on whether a C or an AE lottery is being

evaluated. This method does not assume EUT, and indeed we allow non-EUT specifications. We specify a source-dependent form of utility and probability weighting function and test for violations of ROCL by determining if the subjects evaluate simple and compound lotteries differently.

In both of our methods of evaluation of ROCL, we use data from the 1-in-1 treatment and the 1-in-40 treatment which uses RLIM as the payment protocol. Of course, analysis of the data from the 1-in-40 treatment requires us to assume incentive compatibility with respect to the experiment payment protocol. However, by also analyzing choices from the 1-in-1 treatment we can test if the RLIM itself creates distortions that could be confounded with violations of ROCL. We conclude with discussion of the relative advantages and disadvantages of the econometric tests and the choice pattern tests.

3. Non-Parametric Analysis of Choice Patterns

A. Choice Patterns Where ROCL Predicts Indifference

The basic prediction of ROCL is that subjects who satisfy the axiom are indifferent between a compound lottery and its actuarially-equivalent lottery. We analyze the observed responses from subjects who were presented with any of the 10 pairs that contained both a C lottery and its AE lottery. First, we study the responses from the 32 subjects who were presented with an AE-C pair in 1-in-1 treatment. Then, we study the 620 responses from the 62 subjects who each were presented with all of the 10 AE-C pairs in the 1-in-40 treatment.

We analyze the data separately because, in contrast to the 1-in-40 treatment, any conclusion drawn from the 1-in-1 treatment do not depend on the incentive compatibility of the RLIM. We want to control for the possibility that the observed choice patterns in the 1-in-40 treatment are affected by this payment protocol.¹¹ By analyzing data from the 1-in-1 treatment only, we avoid any possible confounds

¹⁰ These are pairs 31 through 40 of Table B4.

¹¹ An additional consideration is that our interface did not allow expression of indifference, so we test for equal proportions of expressions of strict preference. Even if we had allowed direct expression of

created by the RLIM.

Our null hypothesis is that subjects behave according to ROCL. ROCL predicts that a subject is indifferent between a C lottery and its paired AE lottery, and therefore we should observe equiprobable response proportions between C and AE lotteries in our 10 AE-C pairs. ROCL is violated if, for a given AE-C lottery pair, we observe that the proportion C lottery choices is significantly different from the proportion of AE lottery choices.

We do not find statistical evidence to reject the basic ROCL prediction of indifference in the 1-in-1 treatment, although we do find statistical evidence to support violations of ROCL in the 1-in-40 treatment. Thus, giving many lottery pairs to individuals *and* using the RLIM to select one choice at random for payoff create distortions in the individual decision-making process that can be confounded with violations of ROCL.

Analysis of Data from the 1-in-1 Treatment. We use a generalized version of the Fisher Exact test to jointly test the null hypothesis that the proportion of subjects who chose the C lottery over the AE lottery in each of the AE-C lottery pairs are the same, as well as the Binomial Probability test to evaluate our null hypothesis of equiprobable choice in each of the AE-C lottery pairs.

We do not observe statistically significant violations of the ROCL indifference prediction in the 1-in-1 treatment. Table 2 presents the generalized Fisher Exact test for all AE-C lottery pair choices, and the test's *p*-value of 0.342 provides support for the null hypothesis. We see from this test that the proportions are the same across pairs. We now use a series of Binomial Probability tests to see if the proportions are different from 50%. Table 3 shows the Binomial Probability test applied individually to each of the AE-C lottery pairs for which we have observations. We see no evidence to reject the null

indifference, we have no way of knowing if subjects were in fact indifferent but preferred to use their own randomizing device (in their heads). The same issue confronts tests of mixed strategies in strategic games.

hypothesis that subjects chose the C and the AE lotteries in equal proportions, as all *p*-values are insignificant at any reasonable level of confidence. The results of both of these tests suggest that ROCL is satisfied in the 1-in-1 treatment.

Analysis of Data from the 1-in-40 Treatment The strategy to test the ROCL prediction of indifference in this treatment is different from the one used in the 1-in-1 treatment, given the repeated measures we have for each subject in the 1-in-40 treatment. We now use the Cochran Q test to evaluate whether the proportion of subjects who choose the C lottery is the same in each of the 10 AE-C lottery pairs. A significant difference of proportions identified by this test is sufficient to reject the null prediction of indifference. Of course, an insignificant difference of proportions would require us to additionally verify that the common proportion across pairs the pairs is indeed 50% before we fail to reject the null hypothesis of indifference.

We observe an overall violation of the ROCL indifference prediction in the 1-in-40 treatment. Table 4 reports the results of the Cochran Q test, as well as summary statistics of the information used to conduct the test. The Cochran Q test yields a *p*-value of less than 0.0001, which strongly suggests rejection of the null hypothesis of equiprobable proportions. We conclude, for at least for one of the AE-C lottery pairs, that the proportion of subjects who chose the C lottery is not equal to 50%. This result is a violation of ROCL and we cannot claim that subjects satisfy ROCL and choose at random in all of the 10 AE-C lottery pairs in the 1-in-40 treatment.

¹² The Binomial Probability test is inappropriate in this setting, as it assumes independent observations. Obviously, observations are not independent when each subject makes 40 choices in this treatment.

¹³ For example, suppose there were only 2 AE-C lottery pairs. If the Cochran Q test finds a significant difference, we conclude that the proportion of subjects choosing the C lottery is not the same in the two lottery pairs. Therefore, even if the proportion for one of the pairs was truly equal to 50%, the test result would imply that the other proportion is not statistically equal to 50%, and thus indifference fails.

B. Choice Patterns Where ROCL Predicts Consistent Choices

Suppose a subject is presented with a given S-C lottery pair, and further assume that she prefers the C lottery over the S lottery. If the subject satisfies ROCL and is separately presented with a second pair of lotteries consisting of the same S lottery and the AE lottery of the previously-presented C lottery, then she would prefer and should choose the AE lottery. Similarly, of course, if she instead prefers the S lottery when presented separately with a given S-C lottery pair, then she should choose the S lottery when presented with the corresponding S-AE lottery pair.

Recall that each of the 15 S-C lottery pairs in Table B2 has a corresponding S-AE pair in Table B3. Therefore, we can construct 15 comparisons of lottery pairs that constitute 15 consistency tests of ROCL. In the 1-in-40 treatment we again must assume that the RLIM is incentive compatible, and we again use data from the 1-in-1 treatment to control for possible confounds created by the RLIM. We must now assume homogeneity in risk preferences for the analysis of behavior in the 1-in-1 treatment, since we are making across-subject comparisons. However, in the next section we will present econometric analysis which allows for heterogeneity in risk preferences and test if a violation of the homogeneity assumption is confounded with a violation of ROCL.

Our hypothesis is that a given subject chooses the S lottery when presented with the S-C lottery pair if and only if the same subject also chooses the S lottery when presented with the corresponding S-AE lottery pair. Therefore, ROCL is satisfied if we observe that the proportion of subjects who choose the S lottery when presented with a S-C pair is equal to the proportion of subjects who choose the S lottery when presented with the corresponding S-AE pair. Conversely, ROCL is violated if we observe unequal proportions of choosing the S lottery across a S-C pair and linked S-AE pair.

We do not find evidence to reject the consistency in patterns implied by ROCL in the 1-in-1

¹⁴ Notice that this is equivalent to stating the null hypothesis using the C and AE lotteries. We chose to work with the S lottery for simplicity.

treatment, while we do find evidence of violations of ROCL in the 1-in-40 treatment. As in the case of the ROCL indifference prediction, we conclude that giving many lottery pairs to individuals and using the RLIM to select one choice at random for payoff create distortions in the individual choice making process that can be confounded with violations of ROCL.

Analysis of Data from the 1-in-1 Treatment We use the Cochran-Mantel-Haenszel (CMH) test to test the joint hypothesis that in all of the 15 paired comparisons, subjects choose in the same proportion the S lottery when presented with the S-C lottery pair and its linked S-AE lottery pair. ¹⁵ If the CMH test rejects the null hypothesis, then we interpret this as evidence of overall ROCL-inconsistent observed behavior. We also use the Fisher Exact test to evaluate individually the consistency predicted by ROCL in each of the 15 linked comparisons of S-C pairs and S-AE pairs for which we have enough data to conduct the test.

We do not reject the ROCL consistency prediction. The CMH test does not reject the joint null hypothesis that the proportion of subjects chose the S lottery when they were presented with any given S-C pair is equal to the proportion of subjects that chose the S lottery when they were presented with the corresponding S-AE pair. The χ^2 -statistic for the CMH test with the continuity correction is equal to 2.393 with a corresponding *p*-value of 0.122. Similarly, the Fisher Exact tests presented in Table 5 show only in one comparison the *p*-value is less than 0.05. These results suggest that the ROCL consistency prediction holds in the 1-in-1 treatment. However, as we

¹⁵ The proportion of subjects who choose the S lottery when presented with a S-C pair, or its paired S-AE lottery pair, has to be equal within each paired comparison, but can differ across comparisons. More formally, the CMH test evaluates the null hypothesis that the odds ratio of each of the 15 contingency tables constructed from the 15 paired comparisons are jointly equal to 1.

¹⁶ We follow Li, Simon and Gart [1979] and use the continuity correction to avoid possible misleading conclusions from the test in small samples.

Analysis of Data from the 1-in-40 Treatment We use the Cochran Q test coupled with the Bonferroni-Dunn (B-D) correction procedure¹⁷ to test the hypothesis that subjects choose the S lottery in the same proportion when presented with linked S-C and S-AE lottery pairs. The B-D procedure takes into account repeated comparisons and allows us to maintain a familywise error rate across the 15 paired comparisons of S-C and S-AE lottery pairs.

We find evidence to reject the ROCL consistency prediction. Table 6 shows the results of the B-D method¹⁸ for each of the 15 paired comparisons. Table 6 provides evidence that with a 5% familywise error rate, subjects choose the S lottery in different proportions across linked S-C lottery pairs and S-AE lottery pairs in two comparisons: Pair 1 vs. Pair 16 and Pair 3 vs. Pair 18. This implies that the ROCL prediction of consistency is rejected in 2 of our 15 consistency comparisons.

We are also interested in studying the patterns of violations of ROCL. A pattern inconsistent with ROCL would be subjects choosing the S lottery when presented with a given S-C lottery pair, but switching to prefer the AE lottery when presented with the matched S-AE pair. We construct 2 × 2 contingency tables that show the number of subjects in any given matched pair who exhibit each of the four possible choice patterns: (i) always choosing the S lottery; (ii) choosing the S lottery when presented with a S-C pair and switching to prefer the AE lottery when presented with the matched S-AE pair; (iii) choosing the C lottery when presented with a S-C pair and switching to prefer the S lottery when

The B-D method is a *post-hoc* procedure that is conducted after calculating the Cochran Q test. The first step is to conduct the Cochran Q test to evaluate the null hypothesis that the proportions of individuals who choose the S lottery is the same in all 15 S-C and 15 S-AE linked lottery pairs. If this null is rejected the B-D method involves calculating a critical value d that takes into account all the information of the 30 lottery pairs. The B-D method allows us to test the statistical significance of the observed difference between proportions of subjects who choose the S lottery in any given paired comparison. Define p_1 as the proportion of subjects who choose the S lottery when presented with a given S-AE lottery pair. Similarly, define p_2 as the proportion of subjects who chose the S lottery in the paired S-C lottery pair. The B-D method rejects the null hypothesis that $p_1 = p_2$ if $|p_1 - p_2| > d$. In this case we would conclude that the observed difference is statistically significant. This is a more powerful test than conducting individual tests for each paired comparison because the critical value d takes into account the information of all 15 comparisons. See Sheskin [2004; p. 871] for further details of the B-D method.

¹⁸ The Cochran Q test rejected its statistical null hypothesis χ^2 statistic 448.55, 29 degrees of freedom and *p*-value<0.0001.

presented with the matched S-AE; and (iv) choosing the C lottery when presented with the S-C lottery and preferring the AE lottery when presented with the matched S-AE.

Since we have paired observations, we use the McNemar test to evaluate the null hypothesis of equiprobable occurrences of discordant choice patterns (ii) and (iii) within each set of matched pairs. We find a statistically significant difference in the number of (ii) and (iii) choice patterns within 4 of the 15 matched pairs. Table 7 reports the exact *p*-values for the McNemar test. The McNemar test results in *p*-values less than 0.05 in four comparisons: Pair 1 vs. Pair 16, Pair 3 vs. Pair 18, Pair 10 vs. Pair 25 and Pair 13 vs. Pair 28. Moreover, the odds ratios of the McNemar tests suggest that the predominant switching pattern is choice pattern (iii): subjects tend to switch from the S lottery in the S-AE pair to the C lottery in the S-C pair. The detailed contingency tables for these 4 matched pairs show that the number of choices consistent with pattern (iii) is considerably greater than the number of choices consistent with (ii).

4. Estimated Preferences from Observed Choices

We now estimate preferences from observed choices, and evaluate whether behavior is consistent with ROCL. Additionally, we test for a treatment effect to determine the impact of RLIM on preferences.

A. Econometric Specification

Assume that utility of income is defined by

$$U(x) = x^{(1-r)}/(1-r)$$
 (1)

where x is the lottery prize and $r \ne 1$ is a parameter to be estimated. For r=1 assume $U(x) = \ln(x)$ if needed.

¹⁹ These violations of ROCL are also supported by the B-D procedure if the familywise error rate is set to 10%.

Thus r is the coefficient of CRRA: r=0 corresponds to risk neutrality, r<0 to risk loving, and r>0 to risk aversion. Let there be J possible outcomes in a lottery, and denote outcome $j\in J$ as x_j . Under EUT the probabilities for each outcome x_j , $p(x_j)$, are those that are induced by the experimenter, so expected utility is simply the probability weighted utility of each outcome in each lottery i:

$$EU_{i} = \sum_{i=1,j} [p(x_{i}) \times U(x_{i})].$$
 (2)

The EU for each lottery pair is calculated for a candidate estimate of r, and the index

$$\nabla EU = EU_R - EU_L \tag{3}$$

is calculated, where EU_L is the "left" lottery and EU_R is the "right" lottery as presented to subjects. This latent index, based on latent preferences, is then linked to observed choices using a standard cumulative normal distribution function $\Phi(\nabla EU)$. This "probit" function takes any argument between $\pm \infty$ and transforms it into a number between 0 and 1. Thus we have the probit link function,

$$prob(choose lottery R) = \Phi(\nabla EU)$$
 (4)

Even though this "link function" is common in econometrics texts, it forms the critical statistical link between observed binary choices, the latent structure generating the index ∇EU , and the probability of that index being observed. The index defined by (3) is linked to the observed choices by specifying that the R lottery is chosen when $\Phi(\nabla EU) > \frac{1}{2}$, which is implied by (4).

The likelihood of the observed responses, conditional on the EUT and CRRA specifications being true, depends on the estimates of r given the above statistical specification and the observed choices. The "statistical specification" here includes assuming some functional form for the cumulative density function (CDF). The conditional log-likelihood is then

$$\ln L(\mathbf{r}; \mathbf{y}, \mathbf{X}) = \sum_{i} \left[(\ln \Phi(\nabla EU) \times \mathbf{I}(\mathbf{y}_{i} = 1)) + (\ln (1 - \Phi(\nabla EU)) \times \mathbf{I}(\mathbf{y}_{i} = -1)) \right]$$
 (5)

where $I(\cdot)$ is the indicator function, $y_i = 1(-1)$ denotes the choice of the right (left) lottery in risk aversion task i, and **X** is a vector of individual characteristics reflecting age, sex, race, and so on.

Harrison and Rutström [2008; Appendix F] review procedures that can be used to estimate

structural models of this kind, as well as more complex non-EUT models, with the goal of illustrating how to write explicit maximum likelihood (ML) routines that are specific to different structural choice models. It is a simple matter to correct for multiple responses from the same subject ("clustering"), if needed.

It is also a simple matter to generalize this ML analysis to allow the core parameter r to be a linear function of observable characteristics of the individual or task. We extend the model to be $r = r_0 + R \times X$, where r_0 is a fixed parameter and R is a vector of effects associated with each characteristic in the variable vector X. In effect, the unconditional model assumes $r = r_0$ and estimates r_0 . This extension significantly enhances the attraction of structural ML estimation, particularly for responses pooled over different subjects and treatments, since one can condition estimates on observable characteristics of the task or subject.

In our case we also extend the structural parameter to take on different values for the lotteries presented as compound lotteries. That is, (1) applies to the evaluation of utility for all simple lotteries and a different CRRA risk aversion coefficient r + rc applies to compound lotteries, where rc captures the additive effect of evaluating a compound lottery. Hence, for compound lotteries, the decision maker employs the utility function

$$U(x \mid compound lottery) = x^{(1-r-rc)}/(1-r-rc)$$
 (1')

instead of (1), and we would restate (1) as

$$U(x \mid simple lottery) = x^{(1-r)}/(1-r)$$
 (1")

for completeness. Specifying preferences in this manner provide us with a structural test for ROCL. If rc = 0 then this implies that compound lotteries are evaluated identically to simple lotteries, which is consistent with ROCL. However, if rc ≠ 0, as conjectured by Smith [1969] for objective and subjective compound lotteries, then, decision-makers violate ROCL in a certain source-dependent manner, where

the "source" here is whether the lottery is simple or compound. As stressed by Smith [1969], $rc \neq 0$ for *subjective* lotteries provides a direct explanation for the Ellsberg Paradox, but is much more readily tested on the domain of objective lotteries. Of course, the linear specification r + rc is a parametric convenience, but the obvious one to examine initially.

An important extension of the core model is to allow for subjects to make some *behavioral* errors. The notion of error is one that has already been encountered in the form of the statistical assumption that the probability of choosing a lottery is not 1 when the EU of that lottery exceeds the EU of the other lottery. This assumption is clear in the use of a non-degenerate link function between the latent index ∇ EU and the probability of picking a specific lottery as given in (4). If there were no errors from the perspective of EUT, this function would be a step function: zero for all values of ∇ EU<0, anywhere between 0 and 1 for ∇ EU=0, and 1 for all values of ∇ EU>0.

We employ the error specification originally due to Fechner and popularized by Hey and Orme [1994]. This error specification posits the latent index

$$\nabla EU = (EU_R - EU_I)/\mu \tag{3'}$$

instead of (3), where μ is a structural "noise parameter" used to allow some errors from the perspective of the deterministic EUT model. This is just one of several different types of error story that could be used, and Wilcox [2008] provides a masterful review of the implications of the alternatives.²¹ As $\mu \rightarrow 0$ this specification collapses to the deterministic choice EUT model, where the choice is strictly determined by

²⁰ Abdellaoui, Baillon, Placido and Wakker [2011] conclude that different probability weighting functions are used when subjects face risky processes with known probabilities and uncertain processes with subjective processes. They call this "source dependence," where the notion of a source is relatively easy to identify in the context of an artefactual laboratory experiment, and hence provides the tightest test of this proposition. Harrison [2011] shows that their conclusions are an artefact of estimation procedures that do not take account of sampling errors. A correct statistical analysis that does account for sampling errors provides no evidence for source dependence using their data. Of course, failure to reject a null hypothesis could just be due to samples that are too small.

²¹ Some specifications place the error at the final choice between one lottery or after the subject has decided which one has the higher expected utility; some place the error earlier, on the comparison of preferences leading to the choice; and some place the error even earlier, on the determination of the expected utility of each lottery.

the EU of the two lotteries; but as μ gets larger and larger the choice essentially becomes random. When μ =1 this specification collapses to (3), where the probability of picking one lottery is given by the ratio of the EU of one lottery to the sum of the EU of both lotteries. Thus μ can be viewed as a parameter that flattens out the link functions as it gets larger.

An important contribution to the characterization of behavioral errors is the "contextual error" specification proposed by Wilcox [2011]. It is designed to allow robust inferences about the primitive "more stochastically risk averse than," and posits the latent index

$$\nabla EU = ((EU_R - EU_I)/\nu)/\mu \tag{3"}$$

instead of (3'), where ν is a new, normalizing term for each lottery pair L and R. The normalizing term ν is defined as the maximum utility over all prizes in this lottery pair minus the minimum utility over all prizes in this lottery pair. The value of ν varies, in principle, from lottery choice pair to lottery choice pair: hence it is said to be "contextual." For the Fechner specification, dividing by ν ensures that the normalized EU difference [(EU_R - EU_L)/ ν] remains in the unit interval for each lottery pair. The term ν does not need to be estimated in addition to the utility function parameters and the parameter for the behavioral error term, since it is given by the data and the assumed values of those estimated parameters.

The specification employed here is the source-dependent CRRA utility function from (1') and (1"), the Fechner error specification using contextual utility from (3"), and the link function using the normal CDF from (4). The log-likelihood is then

$$\ln L(\mathbf{r}, \mathbf{rc}, \boldsymbol{\mu}; \mathbf{y}, \mathbf{X}) = \sum_{i} \left[(\ln \Phi(\nabla EU) \times \mathbf{I}(\mathbf{y}_{i} = 1)) + (\ln (1 - \Phi(\nabla EU)) \times \mathbf{I}(\mathbf{y}_{i} = -1)) \right] \tag{5"}$$

and the parameters to be estimated are r, rc and μ given observed data on the binary choices y and the lottery parameters in X.

It is possible to consider more flexible utility functions than the CRRA specification in (1), but that is not essential for present purposes. We do, however, consider extensions of the EUT model to allow for rank-dependent decision-making under Rank-Dependent Utility (RDU) models.

The RDU model extends the EUT model by allowing for decision weights on lottery outcomes. The specification of the utility function is the same parametric specification (1') and (1") considered for source-dependent EUT. To calculate decision weights under RDU one replaces expected utility defined by (2) with RDU

$$RDU_i = \sum_{i=1,1} \left[w(p(M_i)) \times U(M_i) \right] = \sum_{i=1,1} \left[w_i \times U(M_i) \right]$$
 (2')

where

$$w_{i} = \omega(p_{i} + ... + p_{i}) - \omega(p_{i+1} + ... + p_{i})$$
(6a)

for j=1,..., J-1, and

$$\mathbf{w}_{i} = \omega(\mathbf{p}_{i}) \tag{6b}$$

for j=J, with the subscript j ranking outcomes from worst to best, and $\omega(\cdot)$ is some probability weighting function.

We adopt the simple "power" probability weighting function proposed by Quiggin [1982], with curvature parameter γ:

$$\omega(p) = p^{\gamma} \tag{7}$$

So $\gamma \neq 1$ is consistent with a deviation from the conventional EUT representation. Convexity of the probability weighting function is said to reflect "pessimism" and generates, if one assumes for simplicity a linear utility function, a risk premium since $\omega(p) and hence the "RDU EV" weighted by <math>\omega(p)$ instead of p has to be less than the EV weighted by p. The rest of the ML specification for the RDU model is identical to the specification for the EUT model, but with different parameters to estimate.

It is obvious that one can extend the probability weighting specification to be source-dependent, just as we did for the utility function. Hence we extend (7) to be

$$\omega(p \mid \text{compound lottery}) = p^{\gamma + \gamma c}$$
 (7')

for compound lotteries, and

$$\omega(p \mid \text{simple lottery}) = p^{\gamma}$$
 (7")

for simple lotteries. The hypothesis of source-independence, which is consistent with ROCL, in this case is that yc = 0 and rc = 0.

B. Estimates

Analysis of Data from the 1-in-1 Treatment We focus first on the estimates obtained in the 1-in-1 treatment, since this controls for the potentially contaminating effects of the RLIM on our inferences about ROCL. Of course, this requires us to account for subject heterogeneity, and so we control for heterogeneity in risk preferences. We include the effects of allowing for a series of binary demographic variables: **female** is 1 for women, and 0 otherwise; **senior** is 1 for whether that was the current stage of undergraduate education, and 0 otherwise; **white** is 1 based on self-reported ethnic status; and **gpaHI** is 1 for those reporting a cumulative GPA between 3.25 and 4.0 (at least half A's and B's), and 0 otherwise.

The econometric strategy is to estimate our source-dependent version of EUT and RDU separately and compare the model estimates using the tests developed by Vuong [1989] and Clarke [2003][2007] for non-nested, nested and overlapping models.²² This strategy allows us first to choose the model that best describes the data between the two competing models, and then test the chosen model for violations of ROCL.

Controlling for heterogeneity we find that the data are best described by the source-dependent RDU, and conditional on this model there is no evidence of violations of ROCL. Both the Vuong test and the Clarke test provide statistical evidence that our source-dependent version of RDU is the best model to explain the data in the 1-in-1 treatment.²³ Panel A of Table 8 shows the estimates for the source-dependent RDU. A joint test of the coefficient estimates for the covariates and the constant in

The Vuong test is parametric in the sense that it assumes normality to derive the hypothesis test statistic. We also apply the Clarke test which a distribution-free test.

²³ When we control for heterogeneity, the Vuong test statistic is -1.38 in favor of the source-dependent RDU, with a *p*-value of .083. Further, the Clarke test also gives evidence in favor of the source-dependent RDU with a test statistic equal to 56.

the equation for rc results in a *p*-value of 0.59 and a similar test for parameter γc results in a *p*-value of 0.80. Moreover, a joint test of all the covariates and constants in the equations of rc and γc results in a *p*-value equal to 0.72. If we had assumed that subjects behave according to the source-dependent EUT we would have incorrectly concluded that there is evidence of violations of ROCL, from the joint tests of the effect of all covariates in the rc equation which has a *p*-value less than 0.001. This highlights the importance of choosing the preference representation that best characterizes observed choice behavior.

A joint test of all covariates and constant terms, both in the equations for r and γ , results in a p-value less than 0.01. Figure 6 shows the distributions for estimates of the utility parameter r and the probability weighting parameter γ , ²⁴ which have average values 0.79 and 0.33, respectively. This would imply that the typical subject exhibits diminishing marginal returns in the utility function and probability optimism. ²⁵ Figure 6 also shows the distributions for the point estimates for r + rc and γ + γ c. ²⁶

To summarize, behavior in the 1-in-1 treatment is better characterized by RDU instead of EUT, and we do not find evidence of violations of ROCL with the RDU preference representation. We reach a similar conclusion if preference homogeneity is assumed.

Analysis of Data from the 1-in-40 Treatment Controlling for heterogeneity, we again find that the data are best described by our source-dependent version of RDU, and conditional on this model we find evidence of violations of ROCL. Both the Vuong and Clarke tests provide support for the source-

 $^{^{24}}$ The unobserved parameters r and γ are predicted for each subject by using the vector of individual characteristics and the vector estimated parameters that capture the effect of each covariate.

²⁵ These are only descriptive statistics that may not describe in general our subjects' behavior since there is in uncertainty around the *predicted* values of parameters r and γ. However, a series of tests which test, for each subject, the null hypotheses of linear utility (r = 0) and linearity in probabilities ($\gamma = 1$) result, for all subjects, in *p*-values less than 0.01 and less than 0.05, respectively. These tests are constructed using the standard errors around the covariates' coefficients in the equations for parameters r and γ.

Any comparison between the distributions of r + rc and r, but also between $\gamma + \gamma c$ and γ , has to take into account the uncertainty around the distribution fitting process and the significance of the parameter point estimates.

dependent RDU as the best model to explain the data in the 1-in-40 treatment.²⁷ Panel A of Table 9 shows the estimates for this model. A statistical test for the joint null hypothesis that all covariates in the equations for rc and γ c are jointly equal to zero results in a *p*-value less than 0.001, which provides evidence of violations of ROCL. Similarly, the hypothesis that all the covariates in the equations for parameters r and γ are jointly equal to zero also results in a *p*-value less than 0.001. Figure 7 shows the fitted distributions for the point estimates of the utility and probability weighting parameters across subjects in the 1-in-40 treatment. The average predicted values for r, r + rc, γ and γ + γ c are 0.63, 0.71, 0.95 and 0.62, respectively. This would imply that a typical subject displays diminishing marginal returns when evaluating simple and compound lotteries and exhibits more probability optimism when evaluating compound lotteries.²⁸

If we would have assumed that subjects behave according to the source-dependent version of EUT, we would have incorrectly concluded no violation of ROCL. This conclusion derives from a joint test of the effect of all covariates in the rc equation which result in a *p*-value of 0.67. Panel B of Table 9 shows the estimates for the source-dependent EUT model. This highlights, yet again, the importance of choosing an appropriate preference representation that best describes observed choice behavior.

To summarize, behavior in the 1-in-40 treatment is best characterized by the source-dependent RDU model, and we find evidence of violations of ROCL. We reach the same conclusion if preference homogeneity is assumed.

²⁷ The Vuong test statistic is -5.45 in favor of the source-dependent RDU, with a *p*-value less than .001. Further, the Clarke test also gives evidence in favor of the source-dependent RDU with a test statistic equal to 993.

Again, these are only descriptive statistics that are meant to characterize typical behavior. A series of tests for the null hypotheses of r=0 and rc=0 result in *p*-values less than 0.001 for all subjects. Similar tests for the null hypothesis of $\gamma=1$ result in *p*-values greater than 0.05 for 51 out of 62 subjects. Further, tests for the null hypothesis of $\gamma+\gamma c=1$ result in *p*-values less than 0.05 for 37 out of 62 subjects.

5. Conclusions

Our primary goal is to test the Reduction of Compound Lotteries axiom under objective probabilities. Our conclusions are influenced by the experiment payment protocols used and the assumptions about how to characterize risk attitudes.

We do not find violations of ROCL when subjects are presented with one and only one choice that is played for money. However, when individuals are presented with many choices, and the Random Lottery Incentive Mechanism is used to select one choice for payoff, we do find violations of ROCL. These results are obtained whether one uses non-parametric statistics to analyze choice patterns or structural econometrics to estimate preferences.

The econometric analysis provides more information about the structure of individual decision making process. In the context where individuals face only one choice for payoff and no violations of ROCL are found, the preference representation that best characterizes behavior is the Rank-Dependent Utility model. Similarly, when subjects face many choices, behavior is better characterized by our source-dependent version of the RDU model that also accounts for violations of ROCL.

An important methodological conclusion is that the payment protocol used to pay subjects might create distortions of behavior in experimental settings. This is especially important for our purposes since one of the most popular payment protocols assumes ROCL itself. This issue has been studied and documented by Harrison and Swarthout [2012] and Cox, Sadiraj and Schmidt [2011]. Our results provide further evidence that payment protocols can create confounds and therefore affect hypothesis testing about decision making under risk.

Figure 1: Battery of 40 Lotteries Pairs

Probability Coverage




Figure 2: Tree Representation of a Compound Lottery and its Corresponding Actuarially-Equivalent Simple Lottery




Table 1: Experimental Design

Treatment	Subjects	Choices
1. Pay-1-in-1	133	133
2. Pay-1-in-40	62	2480

Figure 3: Choices Over Compound and Actuarially-Equivalent Lotteries




Figure 4: Choices Over Simple and Compound Lotteries




Figure 5: Choices Over Simple and Actuarially-Equivalent Lotteries



Table 2: Generalized Fisher Exact Test on the

Actuarially-Equivalent Lottery vs. Compound Lottery Pairs

Fisher Exact p-value = 0.342

AE-C Lottery Pair	Observed # of choices of AE lotteries	Observed # of choices of C lotteries	Total
31	0	1	1
32	0	3	3
33	2	5	7
36	4	1	5
37	1	1	2
38	2	1	3
39	1	4	5
40	3	3	6
Total	13	19	32

Note: due to the randomization assignment of lottery pairs to subjects, there were no observations for pairs 34 and 35.

Table 3: Binomial Probability Tests on Actuarially-Equivalent Lottery vs. Compound Lottery Pairs

AE-C Lottery Pair	Total # of observations	Observed # of choices of C lotteries	Observed proportion of choices of C lotteries (p)	<i>p</i> -value for H ₀ : p = 0.5
32	3	3	1	0.25
33	7	5	0.714	0.453
36	5	1	0.2	0.375
37	2	1	0.5	1
38	3	1	0.333	1
39	5	4	0.8	0.375
40	6	3	0.5	1

Note: due to the randomization assignment of lottery pairs to subjects there were no observations for pairs 34 and 35 and only 1 observation for pair 31.

Table 4: Cochran Q Test on the Actuarially-Equivalent

Lottery vs. Compound Lottery Pairs

Cochran's χ^2 statistic (9 d.f) = 86.090 p-value < 0.0001

]	Data
AE-C Lottery Pair	Observed # of choices of C lotteries (out of 62 observations)
31	48
32	46
33	34
34	26
35	24
36	18
37	34
38	35
39	49
40	28

Table 5: Fisher Exact Test on Matched Simple-Compound and Simple-Actuarially-Equivalent Pairs

	Total # of subjects in		Proportion of subjects that	Proportion of subjects that	
Comparison	S-AE Pair	chose the S lottery in the S-AE part (π_1)		chose the S lottery in the S-C pair (π_2)	<i>p</i> -value for H_0 : $\pi_1 = \pi_2$
Pair 1 vs. Pair 16	3	2	1	0.5	0.4
Pair 3 vs. Pair 18	6	2	0.5	0	0.464
Pair 5 vs. Pair 20	1	2	0	1	0.333
Pair 6 vs. Pair 21	3	4	0.67	0.5	1
Pair 7 vs. Pair 22	4	9	1	0.56	0.228
Pair 8 vs. Pair 23	3	4	0.33	0.5	1
Pair 9 vs. Pair 24	3	6	0	0.83	0.048
Pair 11 vs. Pair 26	5	9	0.6	0.56	1
Pair 12 vs. Pair 27	5	2	0.8	1	1
Pair 13 vs. Pair 28	4	1	0.5	1	1
Pair 15 vs. Pair 30	3	1	1	0	0.250

Note: due to the randomization assignment of lottery pairs to subjects, the table only shows the Fisher Exact test for 11 S-AE/S-C comparisons for which there are sufficient data to conduct the test.

Table 6: Bonferroni-Dunn Method on Matched Simple-Compound and Simple-Actuarially-Equivalent Pairs

Matching	Proportion of subjects that chose the S lottery in the S-AE pair (<i>p</i> ₁)	Proportion of subjects that chose the S lottery in the S-C pair (\$\nu_2\$)	$ p_1 - p_2 $
Pair 1 vs. Pair 16	0.871	0.387	0.484
Pair 2 vs. Pair 17	0.984	0.952	0.032
Pair 3 vs. Pair 18	0.887	0.629	0.258
Pair 4 vs. Pair 19	0.226	0.210	0.016
Pair 5 vs. Pair 20	0.403	0.290	0.113
Pair 6 vs. Pair 21	0.742	0.661	0.081
Pair 7 vs. Pair 22	0.677	0.548	0.129
Pair 8 vs. Pair 23	0.548	0.548	0.000
Pair 9 vs. Pair 24	0.258	0.306	0.048
Pair 10 vs. Pair 25	0.919	0.774	0.145
Pair 11 vs. Pair 26	0.581	0.613	0.032
Pair 12 vs. Pair 27	0.565	0.645	0.081
Pair 13 vs. Pair 28	0.871	0.726	0.145
Pair 14 vs. Pair 29	0.742	0.677	0.065
Pair 15 vs. Pair 30	0.387	0.419	0.032

Note: the test rejects the null hypothesis of $p_1=p_2$ if $|p_1-p_2| > d$. The calculation of the critical value d requires that one first define *ex ante* a familywise Type I error rate (α_{FW}). For $\alpha_{FW}=10\%$ the corresponding critical value is 0.133, and for $\alpha_{FW}=5\%$ the critical value is 0.159.

Table 7: McNemar Test on Matched Simple-Compound and Simple-Actuarially-Equivalent Pairs

Treatment: 1-in-40

Matching	Exact <i>p</i> -value	Odds Ratio
Pair 1 vs. Pair 16	<0.0001	0.0625
Pair 2 vs. Pair 17	0.625	0.3333
Pair 3 vs. Pair 18	0.0001	0.0588
Pair 4 vs. Pair 19	1.000	0.8571
Pair 5 vs. Pair 20	0.1671	0.4615
Pair 6 vs. Pair 21	0.3323	0.5454
Pair 7 vs. Pair 22	0.1516	0.5000
Pair 8 vs. Pair 23	1.000	1
Pair 9 vs. Pair 24	0.6072	0.6667
Pair 10 vs. Pair 25	0.0352	0.2500
Pair 11 vs. Pair 26	0.8238	1.222
Pair 12 vs. Pair 27	0.4049	1.555
Pair 13 vs. Pair 28	0.0117	0.100
Pair 14 vs. Pair 29	0.5034	0.6667
Pair 15 vs. Pair 30	0.8388	1.1818


Table 8: Estimates of Source-Dependent RDU and EUT Models

Allowing for Heterogeneity

Data from the 1-in-1 treatment (N=133). Estimates of the Fechner error parameter omitted.

Param	eter Covariate	Point Estimate	Standard Error	<i>p</i> -value	95% Confide	ence Interval
			rce-Dependent RD			
	female	0.301	0.077	< 0.001	0.150	0.451
	senior	-0.044	0.060	0.459	-0.161	0.073
r	gpaHI	0.034	0.056	0.552	-0.077	0.144
	white	-0.141	0.095	0.136	-0.326	0.044
	constant	0.636	0.066	< 0.001	0.507	0.764
	female	-0.142	0.089	0.109	-0.316	0.032
	senior	-0.007	0.024	0.784	-0.054	0.041
rc	gpaHI	-0.002	0.013	0.856	-0.027	0.023
	white	-0.073	0.085	0.391	-0.240	0.094
	constant	0.142	0.091	0.118	-0.036	0.319
	female	-0.231	0.199	0.244	-0.621	0.158
	senior	-0.011	0.129	0.932	-0.264	0.242
γ	gpaHI	0.019	0.132	0.888	-0.239	0.276
	white	0.059	0.191	0.757	-0.315	0.433
	constant	0.472	0.130	< 0.001	0.218	0.726
	female	-0.066	0.158	0.675	-0.375	0.243
NC	senior	0.052	0.157	0.741	-0.256	0.360
γс	white	0.061	0.182	0.736	-0.296	0.419
	constant	-0.082	0.119	0.492	-0.314	0.151
		B. Sour	ce-Dependent EUI	(LL=-78.07)		
	female	0.468	0.185	0.011	0.106	0.830
	senior	-0.151	0.125	0.228	-0.396	0.094
r	gpaHI	0.185	0.090	0.039	0.009	0.361
	white	0.068	0.088	0.442	-0.105	0.241
	constant	0.224	0.188	0.233	-0.144	0.593
	female	-0.555	0.208	0.008	-0.963	-0.147
#C	senior	0.148	0.142	0.298	-0.131	0.427
rc	gpaHI	-0.217	0.104	0.036	-0.420	-0.014
	white	-0.097	0.115	0.400	-0.323	0.129
	constant	0.733	0.212	0.001	0.317	1.149

Figure 6: Distribution of Parameter Estimates from the RDU Specification in the 1-in-1 Treatment Assuming Heterogeneity in Preferences







Table 9: Estimates of Source-Dependent RDU and EUT Models


Allowing for Heterogeneity

Data from the 1-in-40 treatment (N=2480). Estimates of the Fechner error parameter omitted.

Parame	eter Covariate	Point Estimate	Standard Error	<i>p</i> -value	95% Confid	ence Interval
		A. Source	-Dependent RDU		35)	
	female	-0.185	0.066	0.005	-0.314	-0.055
	senior	0.035	0.078	0.658	-0.119	0.188
r	gpaHI	0.149	0.096	0.119	-0.039	0.337
	white	-0.233	0.084	0.005	-0.398	-0.069
	constant	0.672	0.072	< 0.001	0.530	0.814
	female	0.084	0.043	0.052	-0.001	0.170
	senior	-0.004	0.049	0.928	-0.101	0.092
rc	gpaHI	-0.071	0.041	0.085	-0.152	0.010
	white	0.060	0.038	0.115	-0.015	0.135
	constant	0.061	0.051	0.232	-0.039	0.161
	female	0.575	0.198	0.004	0.187	0.962
	senior	-0.042	0.221	0.848	-0.476	0.391
γ	gpaHI	-0.380	0.225	0.091	-0.821	0.061
	white	0.312	0.255	0.221	-0.187	0.811
	constant	0.794	0.214	< 0.001	0.374	1.214
	female	-0.311	0.122	0.011	-0.551	-0.071
	senior	0.097	0.126	0.443	-0.150	0.343
γс	gpaHI	0.113	0.134	0.401	-0.150	0.375
	white	-0.230	0.166	0.167	-0.557	0.096
	constant	-0.238	0.130	0.066	-0.493	0.016
		B. Source	-Dependent EUT	(LL= -1504.13	6)	
	female	0.053	0.088	0.547	-0.119	0.225
	senior	-0.014	0.079	0.857	-0.168	0.140
r	gpaHI	-0.041	0.083	0.622	-0.203	0.122
	white	-0.098	0.159	0.538	-0.409	0.213
	constant	0.594	0.104	< 0.001	0.390	0.797
	female	0.007	0.041	0.865	-0.074	0.088
rc	senior	0.019	0.046	0.676	-0.071	0.110
	gpaHI	-0.025	0.047	0.603	-0.117	0.068
	white	0.035	0.037	0.344	-0.038	0.109
	constant	-0.014	0.041	0.727	-0.095	0.066

Figure 7: Distribution of Parameter Estimates from the RDU Specification in the 1-in-40 Treatment Assuming Heterogeneity in Preferences





References

- Abdellaoui, Mohammed; Baillon, Aurélien; Placido, Lætitia and Wakker, Peter P., "The Rich Domain of Uncertainty: Source Functions and Their Experimental Implementation," *American Economic Review*, 101, April 2011, 695-723.
- Andersen, Steffen; Fountain, John; Harrison, Glenn W., and Rutström, E. Elisabet, "Estimating Subjective Probabilities," *Working Paper 2010-06*, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2010.
- Andersen, Steffen; Harrison, Glenn W.; Hole, Arne Rise; Lau, Morten I., and Rutström, E. Elisabet, "Non-Linear Mixed Logit," *Working Paper 2010-07*, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2010; forthcoming, *Theory and Decision*.
- Anscombe, Francis J., and Aumann, Robert .J., "A Definition of Subjective Probability," *Annals of Mathematical Statistics*, 34, 1963, 199-205.
- Beattie, Jane., and Loomes, Graham, "The Impact of Incentives Upon Risky Choice Experiments," *Journal of Risk and Uncertainty*, 14, 1997, 149-162.
- Binswanger, Hans P., "Attitudes Toward Risk: Experimental Measurement in Rural India," *American Journal of Agricultural Economics*, 62, August 1980, 395-407.
- Clarke, Kevin A., "Nonparametric Model Discrimination in International Relations," *Journal of Conflict Resolution*, 47, 2003, 72-93.
- Clarke, Kevin A., "A Distribution-Free Test for Nonnested Model Seleaction," *Political Analysis*, 15(3), 2007, 347-363.
- Cox, James C.; Sadiraj, Vjollca, and Schmidt, Ulrich, "Paradoxes and Mechanism for Choice under Risk," *Working Paper 2011-12*, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2011.
- Cubitt, Robin P.; Starmer, Chris, and Sugden, Robert, "On the Validity of the Random Lottery Incentive System," *Experimental Economics*, 1(2), 1998, 115-131.
- Cubitt, Robin P.; Starmer, Chris, and Sugden, Robert, "Dynamic Choice and Common Ration Effect: An Experimental Investigation," *The Economic Journal*, 108, September 1998a, 1362-1380.
- Ellsberg, Daniel, "Risk, Ambiguity, and the Savage Axioms," *Quarterly Journal of Economics*, 75, 1961, 643-669.
- Ergin, Haluk, and Gul, Faruk, "A Theory of Subjective Compound Lotteries," *Journal of Economic Theory*, 144, 2009, 899-929.
- Fellner, William, "Distortion of Subjective Probabilities as Reaction to Uncertainty," Quarterly Journal

- of Economics, 48(5), November 1961, 670-689.
- Fellner, William, "Slanted Subjective Probabilities and Randomization: Reply to Howard Raiffa and K. R. W. Brewer," *Quarterly Journal of Economics*, 77(4), November 1963, 676-690.
- Galaabaatar, Tsogbadral and Karni, Edi, "Subjective Expected Utility with Incomplete Preferences," Working Paper, Department of Economics, John Hopkins University, 2011.
- Grether, David M., "Testing Bayes Rule and the Representativeness Heuristic: Some Experimental Evidence," *Journal of Economic Behavior & Organization*, 17, 1992, 31-57.
- Halevy, Yoram, "Ellsberg Revisited: An Experimental Study," Econometrica, 75, 2007, 503-536.
- Harrison, Glenn W., "The Rich Domain of Uncertainty: Comment," *Working Paper 2011-13*, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2011.
- Harrison, Glenn W.; Johnson, Eric; McInnes, Melayne M., and Rutström, E. Elisabet, "Measurement With Experimental Controls," in M. Boumans (ed.), *Measurement in Economics:* A Handbook (San Diego, CA: Elsevier, 2007).
- Harrison, Glenn W.; Martínez-Correa, Jimmy, and Swarthout, J. Todd, "Inducing Risk-Neutral Preferences with Binary Lotteries: A Reconsideration," *Draft Working Paper*, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2011.
- Harrison, Glenn W., and Rutström, E. Elisabet, "Risk Aversion in the Laboratory," in J.C. Cox and G.W. Harrison (eds.), *Risk Aversion in Experiments* (Bingley, UK: Emerald, Research in Experimental Economics, Volume 12, 2008).
- Harrison, Glenn W., and Rutström, E. Elisabet, "Expected Utility Theory and Prospect Theory: One Wedding and a Decent Funeral," *Experimental Economics*, 12(2), 2009, 133-158.
- Harrison, Glenn, and Swarthout, J. Todd, "The Independence Axiom and the Bipolar Behaviorist," Working Paper 2012-01, Center for the Economic Analysis of Risk, Robinson College of Business, Georgia State University, 2012.
- Holt, Charles A., and Smith, Angela M., "An Update on Bayesian Updating," *Journal of Economic Behavior & Organization*, 69, 2009, 125-134.
- Karni, Edi, "A Mechanism for Eliciting Probabilities," *Econometrica*, 77(2), March 2009, 603-606.
- Köszegi, Botond, and Rabin, Matthew, "Revealed Mistakes and Revealed Preferences," in A. Caplin and A. Schotter (eds.), *The Foundations of Positive and Normative Economics: A Handbook* (New York: Oxford University Press, 2008).
- Kyburg, Henry E. and Smokler, Howard E., *Studies in Subjective Probability* (New York: Wiley and Sons, 1964).

- Li, Shou-Hua; Simon, Richard M., and Gart, John J., "Small Sample Properties of the Mantel-Haenszel Test," *Biometrika*, 66, 1979, 181-183.
- Loomes, Graham, and Sugden, Robert, "Testing Different Stochastic Specifications of Risky Choice," *Economica*, 65, 1998, 581-598.
- Luce, R. Duncan, and Marley, A.A.J., "On Elements of Chance," *Theory and Decision*, 49, 2000, 97-126.
- Machina, Mark J., and Schmeidler, David, "A More Robust Definition of Subjective Probability," *Econometrica*, 60(4), July 1992, 745-780.
- Machina, Mark J., and Schmeidler, David, "Bayes without Bernoulli: Simple Conditions for Probabilistically Sophisticated Choice," *Journal of Economic Theory*, 67, 1995, 106-128.
- Mantel, Nathan, and Haenszel, William, "Statistical Aspects of the Analysis of Data from Retrospective Studies of Disease," *Journal of the National Cancer Institute*, 22, 1959, 719-748.
- Marascuilo, Leonard A., and Serlin, Ronald C., *Statistical Methods for the Social and Behavioral Sciences* (New York: W. H. Freeman and Company).
- Mathieson, James E., and Winkler, Robert L, "Scoring Rules for Continuous Probability Distributions," *Management Science*, 22(10), June 1976, 1087-1096.
- Oehlert, Gary W., "A Note on the Delta Method," *The American Statistician*, 46(1), February 1992, 27-29.
- Offerman, Theo; Sonnemans, Joep; van de Kuilen, Gijs, and Wakker, Peter P., "A Truth-Serum for Non-Bayesians: Correcting Proper Scoring Rules for Risk Attitudes," *Review of Economic Studies*, 76(4), 2009, 1461-1489.
- Ramsey, Frank P., *The Foundations of Mathematics and Other Logical Essays* (New York: Harcourt Brace and Co, 1926).
- Samuelson, Paul A., "Probability, Utility, and the Independence Axiom," *Econometrica*, 20, 1952, 670-678.
- Savage, Leonard J., "Elicitation of Personal Probabilities and Expectations," *Journal of American Statistical Association*, 66, December 1971, 783-801.
- Savage, Leonard J., *The Foundations of Statistics* (New York: Dover Publications, 1972; Second Edition).
- Schmeidler, David, "Subjective Probability and Expected Utility without Additivity," *Econometrica*, 57, 1989, 571-587.
- Segal, Uzi, "Does the Preference Reversal Phenomenon Necessarily Contradict the Independence Axiom?" *American Economic Review*, 78(1), March 1988, 233-236.

- Segal, Uzi, "Two-Stage Lotteries Without the Reduction Axiom," *Econometrica*, 58(2), March 1990, 349-377.
- Segal, Uzi, "The Independence Axiom Versus the Reduction Axiom: Must We Have Both?" in W. Edwards (ed.), *Utility Theories: Measurements and Applications* (Boston: Kluwer Academic Publishers, 1992).
- Selten, Reinhard; Sadrieh, Abdolkarim, and Abbink, Klaus, "Money Does Not Induce Risk Neutral Behavior, but Binary Lotteries Do Even Worse," *Theory and Decision*, 46, 1999, 211-249.
- Sheskin, David J., Handbook of Parametric and Nonparametric Statistical Procedures (Boca Raton: Chapman & Hall/CRC, 2004; Third Edition).
- Smith, Cedric A.B., "Consistency in Statistical Inference and Decision," *Journal of the Royal Statistical Society*, 23, 1961, 1-25.
- Smith, Vernon L., "Measuring Nonmonetary Utilities in Uncertain Choices: the Ellsberg Urn," *Quarterly Journal of Economics*, 83(2), May 1969, 324-329.
- Starmer, Chris, and Sugden, Robert, "Does the Random-Lottery Incentive System Elicit True Preferences? An Experimental Investigation," *American Economic Review*, 81, 1991, 971-978.
- von Neumann, John. and Morgensten, Oskar, *Theory of Games and Economic Behavior* (Princeton, NJ: Princeton University Press, 1953; Third Edition; Princeton University Paperback Printing, 1980).
- Vuong, Quang H., "Likelihood Ratio Tests for Model Selection and Non-Nested Hypothesis," *Econometrica*, 57(2), March 1989, 307-333.
- Wakker, Peter, Prospect Theory: For Risk and Ambiguity (New York: Cambridge University Press, 2010).
- Wilcox, Nathaniel T., "Stochastic Models for Binary Discrete Choice Under Risk: A Critical Primer and Econometric Comparison," in J. Cox and G.W. Harrison (eds.), Risk Aversion in Experiments (Bingley, UK: Emerald, Research in Experimental Economics, Volume 12, 2008).
- Wilcox, Nathaniel T., "A Comparison of Three Probabilistic Models of Binary Discrete Choice Under Risk," *Working Paper*, Economic Science Institute, Chapman University, March 2010.
- Wilcox, Nathaniel T., "Stochastically More Risk Averse: A Contextual Theory of Stochastic Discrete Choice Under Risk," *Journal of Econometrics*, 162(1), May 2011, 89-104.
- Wooldridge, Jeffrey, "Cluster-Sample Methods in Applied Econometrics," *American Economic Review (Papers & Proceedings)*, 93, May 2003, 133-138.

Appendix A: Instructions (NOT FOR PUBLICATION)


A.1. Instructions for Treatment 1-in-1

Choices Over Risky Prospects

1p

This is a task where you will choose between prospects with varying prizes and chances of winning. You will be presented with one pair of prospects where you will choose one of them. You should choose the prospect you prefer to play. You will actually get the chance to play the prospect you choose, and you will be paid according to the outcome of that prospect, so you should think carefully about which prospect you prefer.

Here is an example of what the computer display of a pair of prospects might look like.




The outcome of the prospects will be determined by the draw of a random number between 1 and 100. Each number between, and including, 1 and 100 is equally likely to occur. In fact, you will be able to draw the number yourself using two 10-sided dice.

In the above example the left prospect pays five dollars (\$5) if the number drawn is between 1 and 40, and pays fifteen dollars (\$15) if the number is between 41 and 100. The blue color in the pie chart corresponds to 40% of the area and illustrates the chances that the number drawn will be

between 1 and 40 and your prize will be \$5. The orange area in the pie chart corresponds to 60% of the area and illustrates the chances that the number drawn will be between 41 and 100 and your prize will be \$15.

Now look at the pie in the chart on the right. It pays five dollars (\$5) if the number drawn is between 1 and 50, ten dollars (\$10) if the number is between 51 and 90, and fifteen dollars (\$15) if the number is between 91 and 100. As with the prospect on the left, the pie slices represent the fraction of the possible numbers which yield each payoff. For example, the size of the \$15 pie slice is 10% of the total pie.

You could also get a pair of prospects in which one of the prospects will give you the chance to play "Double or Nothing." For instance, the right prospect in the following screen image pays "Double or Nothing" if the Green area is selected, which happens if the number drawn is between 51 and 100. The right pie chart indicates that if the number is between 1 and 50 you get \$10. However, if the number is between 51 and 100 a coin will be tossed to determine if you get double the amount. If it comes up Heads you get \$40, otherwise you get nothing. The prizes listed underneath each pie refer to the amounts before any "Double or Nothing" coin toss.




The pair of prospects you choose from is shown on a screen on the computer. On that screen, you should indicate which prospect you prefer to play by clicking on one of the buttons beneath the prospects.

After you have made your choice, raise your hand and an experimenter will come over. It is certain that your one choice will be played out for real. You will roll the two ten-sided dice to determine the outcome of the prospect you chose, and if necessary you will then toss a coin to determine if you get "Double or Nothing."

For instance, suppose you picked the prospect on the left in the last example. If the random number was 37, you would win \$0; if it was 93, you would get \$20.

If you picked the prospect on the right and drew the number 37, you would get \$10; if it was 93, you would have to toss a coin to determine if you get "Double or Nothing." If the coin comes up Heads then you get \$40. However, if it comes up Tails you get nothing from your chosen prospect.

It is also possible that you will be given a prospect in which there is a "Double or Nothing" option no matter what the outcome of the random number. The screen image below illustrates this possibility.



Therefore, your payoff is determined by three things:

- by which prospect you selected, the left or the right;
- by the outcome of that prospect when you roll the two 10-sided dice; and
- by the outcome of a coin toss if the chosen prospect outcome is of the "Double or Nothing" type.

Which prospects you prefer is a matter of personal taste. The people next to you may be presented with a different prospect, and may have different preferences, so their responses should not matter to you. Please work silently, and make your choices by thinking carefully about the prospect you are presented with.

All payoffs are in cash, and are in addition to the \$7.50 show-up fee that you receive just for being here. The only other task today is for you to answer some demographic questions. Your answers to those questions will not affect your payoffs.

Choices Over Risky Prospects

This is a task where you will choose between prospects with varying prizes and chances of winning. You will be presented with a series of pairs of prospects where you will choose one of them. There are 40 pairs in the series. For each pair of prospects, you should choose the prospect you prefer to play. You will actually get the chance to play **one** of the prospects you choose, and you will be paid according to the outcome of that prospect, so you should think carefully about which prospect you prefer.

Here is an example of what the computer display of such a pair of prospects might look like.



The outcome of the prospects will be determined by the draw of a random number between 1 and 100. Each number between, and including, 1 and 100 is equally likely to occur. In fact, you will be able to draw the number yourself using two 10-sided dice.


In the above example the left prospect pays five dollars (\$5) if the number drawn is between 1 and 40, and pays fifteen dollars (\$15) if the number is between 41 and 100. The blue color in the pie chart corresponds to 40% of the area and illustrates the chances that the number drawn will be between 1 and 40 and your prize will be \$5. The orange area in the pie chart corresponds to 60% of the area and illustrates the chances that the number drawn will be between 41 and 100 and your

prize will be \$15.

Now look at the pie in the chart on the right. It pays five dollars (\$5) if the number drawn is between 1 and 50, ten dollars (\$10) if the number is between 51 and 90, and fifteen dollars (\$15) if the number is between 91 and 100. As with the prospect on the left, the pie slices represent the fraction of the possible numbers which yield each payoff. For example, the size of the \$15 pie slice is 10% of the total pie.

Each pair of prospects is shown on a separate screen on the computer. On each screen, you should indicate which prospect you prefer to play by clicking on one of the buttons beneath the prospects.

You could also get a pair of prospects in which one of the prospects will give you the chance to play "Double or Nothing." For instance, the right prospect in the following screen image pays "Double or Nothing" if the Green area is selected, which happens if the number drawn is between 51 and 100. The right pie chart indicates that if the number is between 1 and 50 you get \$10. However, if the number is between 51 and 100 a coin will be tossed to determine if you get double the amount. If it comes up Heads you get \$40, otherwise you get nothing. The prizes listed underneath each pie refer to the amounts before any "Double or Nothing" coin toss.




After you have worked through all of the 40 pairs of prospects, raise your hand and an experimenter will come over. You will then roll two 10-sided dice until a number between 1 and 40 comes up to determine which pair of prospects will be played out. Since there is a chance that any of your 40 choices could be played out for real, you should approach each pair of prospects as if it is the one that you will play out. Finally, you will roll the two ten-sided dice to determine the outcome of the prospect you chose, and if necessary you will then toss a coin to determine if you get "Double or Nothing."

For instance, suppose you picked the prospect on the left in the last example. If the random number was 37, you would win \$0; if it was 93, you would get \$20.

If you picked the prospect on the right and drew the number 37, you would get \$10; if it was 93, you would have to toss a coin to determine if you get "Double or Nothing." If the coin comes up Heads then you get \$40. However, if it comes up Tails you get nothing from your chosen prospect.

It is also possible that you will be given a prospect in which there is a "Double or Nothing" option no matter what the outcome of the random number. The screen image below illustrates this possibility.



Therefore, your payoff is determined by four things:

- by which prospect you selected, the left or the right, for each of these 40 pairs;
- by which prospect pair is chosen to be played out in the series of 40 such pairs using the two 10-sided dice;
- by the outcome of that prospect when you roll the two 10-sided dice; and
- by the outcome of a coin toss if the chosen prospect outcome is of the "Double or Nothing" type.

Which prospects you prefer is a matter of personal taste. The people next to you may be presented with different prospects, and may have different preferences, so their responses should not matter to you. Please work silently, and make your choices by thinking carefully about each prospect.

All payoffs are in cash, and are in addition to the \$7.50 show-up fee that you receive just for being here. The only other task today is for you to answer some demographic questions. Your answers to those questions will not affect your payoffs

Appendix B: Parameters

To construct our battery of 40 lottery pairs, we used several criteria to choose the compound lotteries and their actuarially-equivalent lotteries used in our experiments:

- 1. The lottery compounding task should be as simple as possible. The instructions used by Halevy [2007] are a model in this respect, with careful picture illustrations of the manner in which the stages would be drawn. We wanted to avoid having physical displays, since we had many lotteries. We also wanted to be able to have the computer interface vary the order for us on a between-subject basis, so we opted for a simpler procedure that was as comparable as possible in terms of information as our simple lottery choice interface.
- 2. The lottery pairs should offer reasonable coverage of the Marschak-Machina (MM) triangle and prizes.
- 3. There should be choices/chords that assume parallel indifference curves, as expected under EUT, but the slope of the indifference curve should vary, so that the battery of lotteries can be used to test for a wide range of risk attitudes under the EUT null hypothesis (this criteria was employed for the construction of the basic 69 simple lotteries).
- 4. There should be a number of compound lotteries with their actuarially-equivalent counterparts in the interior of the triangle. Experimental evidence suggests that people tend to comply with the implications of EUT in the interior of the triangle and to violate it on the borders (Conlisk [1989], Camerer [1992], Harless [1992], Gigliotti and Sopher [1993] and Starmer [2000]).
- 5. We were careful to choose lottery pairs with stakes and expected payoff per individual that are comparable to those in the original battery of 69 simple lotteries, since these had been used extensively in other samples from this population.

Our starting point was the battery of 69 lotteries in Table B1 used in Harrison and Swarthout [2012], which in turn were derived from Wilcox [2010]. The lotteries were originally designed in part to satisfy the second and third criteria given above. Our strategy was then to "reverse engineer" the initial lotteries needed to obtain compound lotteries that would yield actuarially-equivalent prospects which already existed in the set of 69 pairs. For instance, the first pair in our battery of 40 lotteries was derived from pair 4 in the battery of 69 (contrast pair 1 in Table B2 with pair 4 in Table B1). We want the distribution of the "risky" lottery in the latter pair to be the actuarially-equivalent prospect of our compound lottery. To achieve this, we have an initial lottery that pays \$10 and \$0 with 50% probability each, and offering "Double or Nothing" if the outcome of the latter prospect is \$10. Hence it offers equal chances of \$20 or \$0 if the DON stage is reached. The \$5 stake was changed to \$0 because DON requires this prize to be among the possible outcomes of the compound lotteries. The actuarially-equivalent lottery of this compound prospect pays \$0 with 75% probability and \$20 with 25% probability, which is precisely the risky lottery in pair 4 of the default battery of 69 pairs. Except for the compound lottery in pair 10 in our set of lotteries, the actuarially-equivalent lotteries play the role of the "risky" lotteries.

Figure B1 shows the coverage of these lottery pairs in terms of the Marschak-Machina triangle. Each prize context defines a different triangle, but the patterns of choice overlap considerably. Figure B1 shows that there are many choices/chords that assume parallel indifference curves, as expected under EUT, but that the slope of the indifference curve can vary, so that the

²⁹ We contemplated using "double or \$5," but this did not have the familiarity of DON.

tests of EUT have reasonable power for a wide range of risk attitudes under the EUT null hypothesis (Loomes and Sugden [1998] and Harrison, Johnson, McInnes and Rutström [2007]). These lotteries also contain a number of pairs in which the "EUT-safe" lottery has a *higher* EV than the "EUT-risky" lottery: this is designed deliberately to evaluate the extent of risk premia deriving from probability pessimism rather than diminishing marginal utility.

The majority of our compound lotteries use a conditional version of the DON device because it allows to obtain good coverage of prizes and probabilities and keeps the compounding representation simple. As noted in the text, one can construct diverse compound lotteries with only two simple components: initial lotteries that either pay two outcomes with 50:50 odds or pay a given stake with certainty, and a conditional DON which pays double a predetermined amount with 50% probability or nothing with equal chance.

In our design, if the subject has to play the DON option she will toss a coin to decide if she gets double the stated amount. One could use randomization devices that allow for probability distributions different from these 50:50 odds, but we want to keep the lottery compounding simple and familiar. Therefore, if one commits to 50:50 odds in the DON option, using exclusively unconditional DON will only allow one to generate compound lotteries with actuarially-equivalent prospects that assign 50% chance to getting nothing. For instance, suppose a compound prospect with an initial lottery that pays positive amounts \$X and \$Y with probability p and p0, respectively, and offers DON for any outcome. The corresponding actuarially-equivalent lottery pays \$2X, \$2Y and \$0 with probabilities p1, p2, p3 and p4, respectively.

The original 69 pairs use 10 contexts defined by three outcomes drawn from \$5, \$10, \$20, \$35 and \$70. For example, the first context consists of prospects defined over prizes \$5, \$10 and \$20, and the tenth context consists of lotteries defined over stakes \$20, \$35 and \$70. As a result of using the DON device, we have to introduce \$0 to the set of stakes from which the contexts are drawn. However, some of the initial lotteries used prizes in contexts different from the ones used for final prizes, so that we could ensure that the stakes for the compounded lottery matched those of the simple lotteries. For example, pair 3 in Table B2 is defined over a context with stakes \$0, \$10 and \$35. The compound lottery of this pair offers an initial lottery that pays \$5 and \$17.50 with 50% chance each and a DON option for any outcome. This allows us to have as final prizes \$0, \$10 and \$35.

Our battery of 40 lotteries uses 6 of the original 10 contexts, but substitute the \$5 stake for \$0. We do not use the other 4 contexts: for them to be distinct from our 6 contexts they would have to have 4 outcomes, the original 3 outcomes plus the \$0 stake required by the DON option. We chose to use only compound lotteries with no more than 3 final outcomes, which in turn requires initial lotteries with no more than 2 outcomes. Accordingly, the initial lotteries of compound prospects are defined over distributions that offer either 50:50 odds of getting any of 2 outcomes or certainty of getting a particular outcome which makes our design simple. It is worth noting that there are compound lotteries composed of initial prospects that offer an amount \$X with 100% probability and a DON option that pays \$2X and \$0 with 50% chance each (see pairs 5, 6 and 14 in Table B2 and pairs 34 and 40 in Table B4). By including this type of "trivial" compound lottery, we provide the basis for ROCL to be tested in its simplest form.

Finally, we included compound lotteries with actuarially-equivalent counterparts in the interior and on the border of the MM triangle, since previous experimental evidence suggests that

this is relevant to test the implications of EUT. Pairs 3, 7, 10, 11, 32, 35 and 38 have compound lotteries with their actuarially-equivalent lotteries in the interior of the triangle.

Additional References

- Camerer, Colin F., "Recent Tests of Generalizations of Expected Utility Theory," in W. Edwards (ed.), *Utility: Theories Measurement, and Applications* (Norwell, MA: Kluwer, 1992).
- Conlisk, John, "Three Variants on the Allais Example," American Economic Review, 79, 1989, 392-407.
- Gigliotti, Gary, and Sopher, Barry, "A Test of Generalized Expected Utility Theory," *Theory and Decision*, 35, 1993, 75-106.
- Harless, David W, "Predictions about Indifference Curves inside the Unit Triangle: a Test of Variants of Expected Utility," *Journal of Economic Behavior and Organization*, 18, 1992, 391-414.
- Starmer, Chris, "D evelopments in Non-Expected Utility Theory: The Hunt for a Descriptive Theory of Choice under Risk," *Journal of Economic Literature*, 38, June 2000, 332-382.

Table B1: Default Simple Lotteries

	Prizes				"Safe"	Lottery Prob	abilities	"Risky" Lottery Probabilities		oabilities		
Pair	Context	Low	Middle	High	Low	Middle	High	Low	Middle	High	EV Safe	EV Risky
1	1	\$5	\$10	\$2 0	0	1	0	0.25	0	0.75	\$10.00	\$16.25
2	1	\$5	\$10	\$2 0	0.25	0.75	0	0.5	0	0.5	\$8.75	\$12.50
3	1	\$5	\$10	\$2 0	0	1	0	0.5	0	0.5	\$10.00	\$12.50
4	1	\$5	\$10	\$2 0	0.5	0.5	0	0.75	0	0.25	\$7.50	\$8.75
5	1	\$5	\$10	\$20	0	1	0	0.25	0.5	0.25	\$10.00	\$11.25
6	1	\$5	\$10	\$20	0.25	0.5	0.25	0.5	0	0.5	\$11.25	\$12.50
7	1	\$5	\$10	\$20	0	0.5	0.5	0.25	0	0.75	\$15.00	\$16.25
8	1	\$5	\$10	\$20	0	0.75	0.25	0.5	0	0.5	\$12.50	\$12.50
9	1	\$5	\$10	\$20	0.25	0.75	0	0.75	0	0.25	\$8.75	\$8.75
10	1	\$5	\$10	\$20	0	1	0	0.75	0	0.25	\$10.00	\$8.75
11	2	\$5	\$10	\$35	0	1	0	0.5	0	0.5	\$10.00	\$20.00
12	2	\$5	\$10	\$35	0	0.75	0.25	0.25	0	0.75	\$16.25	\$27.50
13	2	\$5	\$10	\$35	0.25	0.75	0	0.75	0	0.25	\$8.75	\$12.50
14	2	\$ 5	\$10	\$35	0	0.5	0.5	0.25	0	0.75	\$22.50	\$27.50
15	2	\$5	\$10	\$35	0	0.75	0.25	0.5	0	0.5	\$16.25	\$20.00
16	2	\$5	\$10	\$35	0	1	0	0.75	0	0.25	\$10.00	\$12.50
17	3	\$5	\$10	\$70	0.25	0.75	0	0.5	0	0.5	\$8.75	\$37.50
18	3	\$5	\$10	\$70	0	1	0	0.5	0	0.5	\$10.00	\$37.50
19	3	\$5	\$10	\$70	0.5	0.5	0	0.75	0	0.25	\$7.50	\$21.25
20	3	\$5	\$10	\$70	0	1	0	0.75	0	0.25	\$10.00	\$21.25
21	4	\$5	\$20	\$35	0	1	0	0.25	0	0.75	\$20.00	\$27.50
22	4	\$5	\$20	\$35	0	0.75	0.25	0.25	0	0.75	\$23.75	\$27.50
23	4	\$5	\$20	\$35	0	0.5	0.5	0.25	0	0.75	\$27.50	\$27.50
24	4	\$5	\$20	\$35	0	1	0	0.5	0	0.5	\$20.00	\$20.00
25	4	\$5	\$20	\$35	0.5	0.5	0	0.75	0	0.25	\$12.50	\$12.50

26	4	\$5	\$20	\$35	0	0.75	0.25	0.5	0	0.5	\$23.75	\$20.00
27	4	\$5	\$20	\$35	0.25	0.75	0	0.75	0	0.25	\$16.25	\$12.50
28	5	\$5	\$2 0	\$ 70	0.25	0.75	0	0.5	0	0.5	\$16.25	\$37.50
29	5	\$5	\$20	\$70	0	0.75	0.25	0.25	0	0.75	\$32.50	\$53.75
30	5	\$5	\$2 0	\$ 70	0.5	0.5	0	0.75	0	0.25	\$12.50	\$21.25
31	5	\$5	\$2 0	\$ 70	0.25	0.5	0.25	0.5	0	0.5	\$28.75	\$37.50
32	5	\$5	\$2 0	\$ 70	0.25	0.75	0	0.75	0	0.25	\$16.25	\$21.25
33	5	\$5	\$2 0	\$ 70	0	0.5	0.5	0.25	0	0.75	\$45.00	\$53.75
34	6	\$5	\$35	\$70	0	1	0	0.25	0	0.75	\$35.00	\$53.75
35	6	\$5	\$35	\$70	0.25	0.75	0	0.5	0	0.5	\$27.50	\$37.50
36	6	\$5	\$35	\$70	0	0.75	0.25	0.25	0	0.75	\$43.75	\$53.75
37	6	\$5	\$35	\$70	0.5	0.5	0	0.75	0	0.25	\$20.00	\$21.25
38	6	\$5	\$35	\$70	0	0.5	0.5	0.25	0	0.75	\$52.50	\$53.75
39	6	\$5	\$35	\$70	0	0.75	0.25	0.5	0	0.5	\$43.75	\$37.50
40	6	\$5	\$35	\$70	0.25	0.75	0	0.75	0	0.25	\$27.50	\$21.25
41	6	\$5	\$35	\$70	0	1	0	0.75	0	0.25	\$35.00	\$21.25
			W									
42	7	\$10	\$20	\$35	0	1	0	0.25	0	0.75	\$20.00	\$28.75
43	7	\$10	\$20	\$35	0.25	0.75	0	0.5	0	0.5	\$17.50	\$22.50
44	7	\$10	\$20	\$35	0	1	0	0.25	0.25	0.5	\$20.00	\$25.00
45	7	\$10	\$20	\$35	0	1	0	0.5	0	0.5	\$20.00	\$22.50
46	7	\$10	\$20	\$35	0	1	0	0.25	0.5	0.25	\$20.00	\$21.25
47	7	\$10	\$20	\$35	0	0.75	0.25	0.5	0	0.5	\$23.75	\$22.50
48	7	\$10	\$20	\$35	0	1	0	0.5	0.25	0.25	\$20.00	\$18.75
49	7	\$10	\$20	\$35	0.25	0.75	0	0.75	0	0.25	\$17.50	\$16.25
50	7	\$10	\$20	\$35	0	1	0	0.75	0	0.25	\$20.00	\$16.25
		***	**			. ==			•	o =	***	* 40.00
51	8	\$10	\$20	\$ 70	0.25	0.75	0	0.5	0	0.5	\$17.50	\$40.00
52	8	\$10	\$20	\$ 70	0.5	0.5	0	0.75	0	0.25	\$15.00	\$25.00
53	8	\$10	\$20	\$70	0.25	0.75	0	0.75	0	0.25	\$17.50	\$25.00

54	9	\$10	\$35	\$70	0	1	0	0.25	0	0.75	\$35.00	\$55.00
55	9	\$10	\$35	\$70	0.25	0.75	0	0.5	0	0.5	\$28.75	\$40.00
56	9	\$10	\$35	\$70	0	0.5	0.5	0.25	0	0.75	\$52.50	\$55.00
57	9	\$10	\$35	\$70	0	0.75	0.25	0.5	0	0.5	\$43.75	\$40.00
58	10	\$20	\$35	\$70	0	1	0	0.25	0	0.75	\$35.00	\$57.50
59	10	\$20	\$35	\$70	0.25	0.75	0	0.5	0	0.5	\$31.25	\$45.00
60	10	\$20	\$35	\$70	0	0.75	0.25	0.25	0	0.75	\$43.75	\$57.50
61	10	\$20	\$35	\$70	0	1	0	0.5	0	0.5	\$35.00	\$45.00
62	10	\$20	\$35	\$70	0.5	0.5	0	0.75	0	0.25	\$27.50	\$32.50
63	10	\$20	\$35	\$70	0	1	0	0.25	0.5	0.25	\$35.00	\$40.00
64	10	\$20	\$35	\$70	0.25	0.5	0.25	0.5	0	0.5	\$40.00	\$45.00
65	10	\$20	\$35	\$70	0	0.5	0.5	0.25	0	0.75	\$52.50	\$57.50
66	10	\$20	\$35	\$70	0	1	0	0.5	0.25	0.25	\$35.00	\$36.25
67	10	\$20	\$35	\$70	0.25	0.75	0	0.75	0	0.25	\$31.25	\$32.50
68	10	\$20	\$35	\$70	0	0.75	0.25	0.5	0	0.5	\$43.75	\$45.00
69	10	\$20	\$35	\$70	0	1	0	0.75	0	0.25	\$35.00	\$32.50

Figure B1: Default Simple Lotteries




Table B2: Simple Lotteries vs. Compound Lotteries (Pairs 1-15)

											Com	pound Lott	erv			
					Si	mple Lotte	ry	1	nitial Lotte	ry	I	nitial Lotter	y			
		Final Pri	zes		1	Probabilitie	S		Prizes			Probabilities	s	"Double or Nothing"	EV	EV
Pair	Context	Low	Middle	High	Low	Middle	High	Low	Middle	High	Low	Middle	High	option	Simple	Compound
1	1	\$0	\$10	\$20	0.5	0.5	0	\$ 0	\$10	\$20	0.5	0.5	0	DON if middle	\$5.00	\$5.00
2	1	\$0	\$10	\$20	0	1	0	\$ 0	\$10	\$20	0.5	0.5	0	DON if middle	\$10.00	\$5.00
3 4	2 2	\$0 \$0	\$10 \$10	\$35 \$35	0 0.25	1 0.75	0	\$0 \$0	\$5 \$17.50	\$17.50 \$35	0 0.5	0.5 0.5	0.5	DON for any outcome DON if middle	\$10.00 \$7.50	\$11.25 \$8.75
7	2	90	\$10	933	0.23	0.75	Ü	30	\$17.50	\$33	0.5	0.5	Ü	DON II IIIIddie	97.50	φ0.73
5	3	\$0	\$10	\$ 70	0.25	0.75	0	\$ 0	\$35	\$70	0	1	0	DON for any outcome	\$7.50	\$35.00
6	3	\$0	\$10	\$70	0	1	0	\$0	\$35	\$70	0	1	0	DON for any outcome	\$10.00	\$35.00
7	4	\$0	\$20	\$35	0	1	0	\$0	\$10	\$35	0	0.5	0.5	DON if middle	\$20.00	\$22.50
8	4	\$0	\$20	\$35	0	0.75	0.25	\$ 0	\$17.50	\$35	0	0.5	0.5	DON if middle	\$23.75	\$8.75
9	5	\$0	\$20	\$70	0	0.5	0.5	\$0	\$35	\$70	0	0.5	0.5	DON if middle	\$45.00	\$52.50
10	5	\$0	\$20	\$70	0.5	0	0.5	\$0	\$20	\$35	0	0.5	0.5	DON if high	\$35.00	\$27.50
11	5	\$0	\$20	\$70	0	1	0	\$ 0	\$20	\$35	0	0.5	0.5	DON if high	\$20.00	\$27.50
12	5	\$0	\$20	\$ 70	0	0.75	0.25	\$0	\$35	\$70	0	0.5	0.5	DON if middle	\$32.50	\$52.50
13	6	\$0	\$35	\$70	0	1	0	\$ 0	\$35	\$70	0	0.5	0.5	DON if middle	\$35.00	\$52.50
14	6	\$0	\$35	\$70	0	0.75	0.25	\$0	\$35	\$70	0	1	0	DON for any outcome	\$43.75	\$35.00
15	6	\$0	\$35	\$70	0	0.75	0.25	\$0	\$35	\$ 70	0	0.5	0.5	DON if middle	\$43.75	\$52.50

Table B3: Simple Lotteries vs. Actuarially-Equivalent Lotteries (Pairs 16-30)

		Final P	rizes			imple Lotter	•		arially-Equi		EV Simple	EV Actuarially-
Pair	Context	Low	Middle	High	Low	Middle	High	Low	Middle	High		Equivalent
16	1	\$ 0	\$ 10	\$20	0.5	0.5	0	0.75	0	0.25	\$5.00	\$5.00
17	1	\$0	\$10	\$20	0	1	0	0.75	0	0.25	\$10.00	\$5.00
18	2	\$0	\$10	\$35	0	1	0	0.5	0.25	0.25	\$10.00	\$11.25
19	2	\$ 0	\$10	\$35	0.25	0.75	0	0.75	0	0.25	\$7.50	\$8.75
20	3	\$ 0	\$10	\$70	0.25	0.75	0	0.5	0	0.5	\$7.50	\$35.00
21	3	\$ 0	\$10	\$70	0	1	0	0.5	0	0.5	\$10.00	\$35.00
22	4	\$ 0	\$20	\$35	0	1	0	0.25	0.25	0.5	\$20.00	\$22.50
23	4	\$ 0	\$20	\$35	0	0.75	0.25	0.25	0	0.75	\$23.75	\$8.75
24	5	\$ 0	\$20	\$70	0	0.5	0.5	0.25	0	0.75	\$45.00	\$52.50
25	5	\$ 0	\$20	\$70	0.5	0	0.5	0.25	0.5	0.25	\$35.00	\$27.50
26	5	\$ 0	\$20	\$70	0	1	0	0.25	0.5	0.25	\$20.00	\$27.50
27	5	\$ 0	\$20	\$70	0	0.75	0.25	0.25	0	0.75	\$32.50	\$52.50
28	6	\$ 0	\$35	\$70	0	1	0	0.25	0	0.75	\$35.00	\$52.50
29	6	\$0	\$35	\$70	0	0.75	0.25	0.5	0	0.5	\$43.75	\$35.00
30	6	\$ 0	\$35	\$70	0	0.75	0.25	0.25	0	0.75	\$43.75	\$52.50

Table B4: Actuarially-Equivalent Lotteries vs. Compound Lotteries (Pairs 31-40)

											Com	pound Lott	e r v			
					Actu	arially-Equi	valent	I	Initial Lotte	ry		nitial Lotter			EV	EV
		Final Priz	zes		Lot	tery Probabi	ilities		Prizes		:	Probabilities	3	"Double or Nothing"	Actuarially-	Compound
Pair	Context	Low	Middle	High	Low	Middle	High	Low	Middle	High	Low	Middle	High	option	Equivalent	
31	1	\$0	\$10	\$20	0.75	0	0.25	\$0	\$ 10	\$20	0.5	0.5	0	DON if middle	\$5.00	\$5.00
32 33	2 2	\$0 \$0	\$10 \$10	\$35 \$35	0.5 0.75	0.25 0	0.25 0.25	\$0 \$0	\$5 \$17.50	\$17.50 \$35	0 0.5	0.5 0.5	0.5	DON for any outcome DON if middle	\$11.25 \$8.75	\$11.25 \$8.75
34	3	\$0	\$ 10	\$ 70	0.5	0	0.5	\$ 0	\$35	\$70	0	1	0	DON for any outcome	\$35.00	\$35.00
35	4	\$0	\$20	\$35	0.25	0.25	0.5	\$ 0	\$ 10	\$35	0	0.5	0.5	DON if middle	\$22.50	\$22.50
36	4	\$0	\$20	\$35	0.25	0	0.75	\$0	\$17.50	\$35	0	0.5	0.5	DON if middle	\$8.75	\$8.75
37	5	\$0	\$20	\$70	0.25	0	0.75	\$0	\$35	\$70	0	0.5	0.5	DON if middle	\$52.50	\$52.50
48	5	\$0	\$20	\$70	0.25	0.5	0.25	\$0	\$20	\$35	0	0.5	0.5	DON if high	\$27.50	\$27.50
30	6	\$0	\$35	\$70	0.25	0	0.75	\$0	\$35	\$70	0	0.5	0.5	DON if middle	\$52.50	\$52.50
40	6	\$0	\$35	\$70	0.5	0	0.5	\$0	\$35	\$70	0	1	0	DON for any outcome	\$35.00	\$35.00

Appendix C: Related Literature (NOT FOR PUBLICATION)

Cubitt, Starmer and Sugden [1998a] studied the Reduction of Compound Lotteries Axiom [ROCL] in a 1-in-1 design that gave each subject one and only one problem for real stakes and was conceived to test principles of dynamic choice. Also, Starmer and Sugden [1991], Beattie and Loomes [1997] and Cubitt, Starmer and Sugden [1998] have studied the Random Lottery Incentive Method (RLIM), and as a by-product have tested the ROCL axiom. We focus on the results related to the ROCL axiom: Harrison and Swarthout [2011] review the results related to RLIM.

Cubitt, Starmer and Sugden [1998a] gave to one group of subjects one problem that involved compound lotteries and gave to another group the reduced compound version of the same problem. If ROCL is satisfied, one should see the same pattern of choice in both groups. They cannot find statistically significant violations of ROCL in their design.

Starmer and Sugden [1991] gave their subjects two pairs of lotteries that were designed to test "common consequence" violations of EUT. In each pair i there was a risky (Ri) option and a safe (Si) option. They recruited 160 subjects that were divided into four groups of equal number. Two groups faced one of the two pairs in 1-in-1 treatments, while the other two groups were given both pairs to make a choice over using the RLIM to choose the pair for final payoff. We focus on the latter two groups since RLIM induces four possible compound lotteries: i) (0.5, R1; 0.5, R2), ii) (0.5, R1; 0.5, S2), iii) (0.5, R2; 0.5, S1) and iv) (0.5, S1; 0.5, S2). The lottery parameters were chosen to make compound lotteries ii) and iii) have equal actuarially-equivalent prospects.

They hypothesize that if a *reduction principle* holds, and if any of the induced compound lotteries ii) and iii) above is preferred by a subject, then the other one must be preferred.³⁰ The rejection of this hypothesis is a violation of ROCL, since this axiom implies that two compound lotteries with the same actuarially-equivalent prospects should be equally preferred. Therefore, the null hypothesis in Starmer and Sugden [1991; p. 976] is that "the choice between these two responses to be made at random; as a result, these responses should have the same expected frequency." From the 80 subjects that faced the 1-in-2 treatments, 32.5% of the individuals chose (0.5, R1; 0.5, S2) and 15% chose (0.5, R2; 0.5, S1), thus Starmer and Sugden reject the null hypothesis of equal frequency in choices based on a one-tail test with a binomial distribution and *p*-value=0.017. This pattern is very similar in each of the 1-in-2 treatments; in one of them the proportions are 30% and 15%, whereas in the other they are 35% and 15%. A two-sided Fisher Exact test yields a *p*-value of 0.934, which suggest that these patterns of choices are very similar in both 1-in-2 treatments. Therefore, there is no statistical evidence to support ROCL in their experiment.

Beattie and Loomes [1997] examined 4 lottery choice tasks. The first 3 tasks involved a binary choice between two lotteries, and the fourth task involved the subject selecting one of four possible lotteries, two of which were compound lotteries.³¹ They recruited 289 subjects that were

³⁰ Following Holt [1986], Starmer and Sugden [1991, p. 972] define the reduction principle to be when "compound lotteries are reduced to simple ones by the calculus of probabilities and that choices are determined by the subject's preferences over such reduced lotteries." This is what we call ROCL, in *addition* to some axioms that are maintained for present purpose.

³¹ Beattie and Loomes [1997] use nine prospects: A= (0.2, £15; 0.8, £0), B= (0.25, £10; 0.75, £0), C= (0.8, £0; 0.2, £30), D= (0.8, £5; 0.2, £0), E=(0.8, £15; 0.2, £0), F = (1, £10), G=(1, £4), H=(0.5,£10; 0.5,£0),

randomly assigned to six groups. The first group faced a hypothetical treatment and was paid a flat fee for completing all four tasks. The second group was given a 1-in-4 treatment, and each of the other four groups faced one of the four tasks in 1-in-1 treatments. Sample sizes were 49 for the hypothetical treatment and the 1-in-4 treatment, and a total of 191 in the four 1-in-1 treatments.

Beattie and Loomes [1997; p. 164] find that "there is no support for the idea that two problems involving the same 'reduced form' alternatives – and therefore involving the same difference between expected values – will be treated equivalently." On this basis, their Question 3 in the 1-in-4 treatment would be actuarially-equivalent to their Question 1 in the 1-in-1 treatment. They found that the pattern of choices in both treatments are so different "that a chi-square test rejects with a very great confidence (p<.001) the hypothesis that they are treated equivalently" (p. 164). The p-value<0.001 of the Fisher Exact test provides further support for this violation of ROCL.

Their Question 4 is a task that is similar to the method developed by Binswanger [1980]: subjects are offered an ordered set of choices that increase the average payoff while increasing variance. The difference with the Binswanger procedure is that two of the four choices involved compound lotteries: one paid a given amount of money if two Heads in a row were flipped, and the other paid a higher amount if three Heads in a row were flipped. For responses in Question 4, Beattie and Loomes [1997; p.162]

...conjecture that the REAL [1-in-1] treatment might stimulate the greatest effort to picture the full sequential process [of coin flipping in the compound prospects] and, as a part of that, to anticipate feelings at each stage in the sequence; whereas the HYPO [hypothetical] treatment would be most conducive to thinking of the alternatives in their reduced form as a set of simple lotteries... The RPSP [1-in-4 treatment] might then, both formally and psychologically, represent an intermediate position, making the process less readily imaginable by adding a further stage (the random selection of the problem) to the beginning of the sequence, and reducing but not eliminating the incentive to expend the necessary imaginative effort.

On this basis, they predict that, when answering Question 4, subjects in the hypothetical treatment are more likely to think in *reduced form probability distributions*. Beattie and Loomes consider that this might enhance the salience of the high-payoff option, and thus the compound lotteries are expected to be chosen more frequently in the hypothetical treatment than in the 1-in-1 and 1-in-4 treatments.

Beattie and Loomes [1997; p.165] found support for these conjectures: Their subjects tend to choose the compound lotteries more often in the hypothetical treatment than in the ones with economic incentives (i.e., 1-in-1 and 1-in-4 treatments). They found that under the hypothetical treatment more than 1 in 3 of the sample opted for the compound lotteries; this proportion was reduced in the 1-in-4 treatment to just over 1 in 5; and in the 1-in-1 treatment the proportion fell to 1 in 12. A chi-square test rejects (p-value < 0.01) the hypothesis that there is no difference in

I=(£25 if two Heads in a row are flipped; otherwise nothing) and J=(£62.50 if three Heads in a row are flipped; otherwise nothing). Questions 1 through 3 are binary choices that offer, respectively, A or B, C or D and E or F. In Question 4, the subject must choose the prospect that she prefers the most among G, H, I or J. Options I and J are compound lotteries with 2 and 3 stages, respectively.

patterns across treatments. The Fisher Exact test is consistent with this result.³²

Cubitt, Starmer and Sugden [1998] use common consequence and common ratio pairs of pairs in three experiments. We focus in the first two since the third experiment has no treatments relevant to test ROCL. In the first experiment they compare 1-in-1 choices with 1-in-2 choices. Their comparison rests on subjects not having extreme risk-loving preferences over the pairs of lotteries in the 1-in-2 treatment designed to capture this behavior. Given that this *a priori* assumption is true, and it is generally supported by their data, the lottery pairs in each of the 1-in-2 treatments were chosen to generate compound prospects with actuarially-equivalent lotteries equal to the prospects in each of the 1-in-1 treatments.

If ROCL is satisfied, the distribution of responses between risky and safe lotteries should be the same in both treatments. The *p*-value from the Fisher Exact test in one of the 1-in-1 and 1-in-2 treatment comparisons³³ is 0.14, which suggests that ROCL is most likely violated.³⁴

Similarly, in the second experiment the 1-in-2 treatment induced compound lotteries with actuarially-equivalent prospects equal to the lottery choices in one of their 1-in-1 treatment. In the latter, 52% of the 46 subjects chose the risky lottery, whereas 38% of the 53 subjects in the 1-in-2 treatment chose the risky prospect. These choice patterns suggest that ROCL does not hold in the second experiment.³⁵

 $^{^{32}}$ We test the similarity between treatments of the proportions of subjects that chose each of the four prospects in Question 4. The two-sided Fisher Exact test applied to the hypothetical and the 1-in-1 treatments rejects the hypothesis of no difference in choice patterns (*p*-value = 0.001). The *p*-value for the comparison of the same four choices between the hypothetical and the 1-in-4 treatments is 0.473.

³³ Groups 1.1 and 1.3 in their notation.

³⁴ The proportions of subjects that chose the risky prospect in the other 1-in-1 and 1-in-2 treatments (groups 1.2 and 1.4 in their notation) are close: 50% and 55%, respectively. However, we cannot perform the Fisher Exact test for this 1-in-1 and 1-in-2 comparison, since the compound lotteries induced by the 1-in-2 treatment have actuarially-equivalent prospects equal to the ones in the 1-in-1 treatment only if the subjects do not exhibit extreme risk-loving preferences. Since 8% of the subjects in this 1-in-2 treatment exhibited risk-loving preferences, one cannot perform the Fisher test because this contaminates the comparison between the compound lotteries and their actuarially-equivalent counterparts.

³⁵ Since one of the subjects in the 1-in-2 treatment (group 2.3) exhibited risk-loving preferences, we cannot perform the Fisher Exact test for the reasons explained earlier.

Appendix D: Nonparametric Tests (NOT FOR PUBLICATION)

A. Choice Patterns Where ROCL Predicts Indifference

Research Hypothesis: Subjects are indifferent between a compound (C) lottery and its paired actuarially-equivalent (AE) lottery, and therefore the choice between both lotteries is made at random in any of our 10 AE-C pairs in Table B4. As a result we should observe equiprobable response proportions between C and AE lotteries. ROCL is rejected if we can provide statistical evidence that the proportion of observations in which subjects chose a C lottery over its AE lottery is different from the proportion of choices in which subjects chose the AE lottery over the C lottery.

Structure of data sets: We analyze the observed responses from subjects who were presented with any of the 10 pairs described in Table B4 and which contain both a C lottery and its AE lottery. First, we study the responses from 32 subjects who were presented with one and only one of the AE-C pairs in 1-in-1 treatment. We also study the 620 responses from the 62 subjects that were presented with all of the 10 AE-C pairs in the 1-in-40 treatment. In terms of the statistical literature, the responses to each of the AE-C pairs in the 1-in-1 constitute an "independent" sample. This means that subjects are presented with one and only one choice, thus one observation does not affect any other observation in the sample. Conversely, the responses to the AE-C pairs in the 1-in-40 constitute 10 "dependent" samples since each of the 62 subjects responded to each of the 10 AE-C pairs.

We analyze the data separately because, in contrast to the 1-in-40 treatment which uses the random lottery incentive mechanism (RLIM) as payment protocol, any conclusion drawn from the 1-in-1 treatment does not depend in the independence axiom assumed by the RLIM. We want to control for the possibility that the observed choice patterns in the 1-in-40 treatment are affected by this payment protocol. This means that any failure to see indifference between a C lottery and its and AE lottery in our data could be explained by confounds created by the payment protocol. By analyzing data from the 1-in-1 treatment only, we avoid possible confounds created by the RLIM.

1-in-1 Treatment

We apply the Binomial probability test to each of the AE-C pair for which there is sufficient data to conduct the test. The latter allows us to test individually for each AE-C pair if subjects choose the C lottery and the AE lottery in equal proportions. We also use a generalized version of the Fisher Exact test that allows us to jointly test the statistical null hypothesis that the proportions of subjects that chose the C lottery over the AE lottery in each of the AE-C lottery pairs are the same. Both test can provide statistical evidence of the overall performance of the ROCL indifference prediction.

Statistical Null Hypothesis of the Binomial Probability Test: For a given AE-C lottery pair, the proportion of subjects that choose the C lottery is 50 %. This is equivalent to test the claim that subjects choose the AE lottery and the C lottery in equal proportions when they are presented with a given AE-C lotteries.

If this statistical null hypothesis is not rejected then we conclude that there is evidence to support the claim that, for a given AE-C pair, subjects choose the AE and the C lotteries in equal

proportions. If the null hypothesis of the test is rejected, then we concluded that subjects choose the AE and the C lotteries in different proportions. The rejection (acceptance) of the statistical null hypothesis implies that the research hypothesis is rejected (accepted) and we conclude that there is evidence to support the claim that the basic ROCL indifference prediction is violated (satisfied) in a given AE-C pair.

This is an appropriate test to use for this treatment because it allows us to test the equality of proportions implied by the ROCL prediction. More importantly, the basic assumptions of the test are satisfied by the data. As described by Sheskin[2004; p. 245], the assumptions are: (i) each of the *n* observations are independent (i.e., the outcome of one observation is not affected by the outcome of another observation), (ii) each observation is randomly selected from a population, and (iii) each observation can be classified only two into mutually exclusive categories. Assumption (i) is satisfied because each subject makes one and only one choice. Assumption (ii) is also satisfied because each subject is randomly recruited from the extensive subject pool of the EXCEN laboratory at Georgia State University. Finally (iii) is also satisfied because subjects can only choose either the C lottery or the AE lottery in each of the 10 AE-C pairs.

Statistical Null Hypothesis of the Generalized Fisher Exact Test: The proportion of individuals choosing the C lotteries is the same for all the AE-C lottery pairs.

The contingency table used in the test has the following structure:

	Cho		
Pair	AE Lottery	C Lottery	Total
AE-C pair 1	n11	n12	n11 + n12
AE-C pair 2	n21	n22	n21 + n22
	•	•	·
AE-C pair r	nr1	nr2	nr1 + nr2
Total	Σi-1,2,,r (ni1)	Σi-1,2,,r (ni2)	$\Sigma_{i-1,2,,r}$ (ni1+ni2)

The number in each cell is defined as follows. The symbol n11 represents the number of individuals that chose the AE lottery when they were presented with the AE-C lottery pair 1. The symbol n12 represents the number of individuals that chose the C lottery when they were presented with the same AE-C pair 1. The sum n11+n12 represents the total number of subjects that were presented with the AE-C lottery pair 1. The interpretation of ni1 and ni2, for i=2, 3, ..., r, can be similarly derived. The generalized Fisher Exact test tests the null hypothesis that the proportion of subjects that choose the C lottery is statistically the same for all of the r AE-C lottery pairs used in the table. Formally, the statistical null hypothesis is Ho: p1 = p2 = ... = pr, where pi = ni2 / (ni1 + ni2) for i=1, 2, 3, ..., r.

We use this test in conjunction with the Binomial probability test applied individually to each of the AE-C lottery pairs to make stronger claims of the overall performance of ROCL. If the

Binomial probability test does not reject its statistical null hypothesis for each of the AE-C lottery pairs and if the statistical null hypothesis of the generalized Fisher Exact test is not rejected, we can conclude that the proportion of subjects that chose the C lottery is statistically the same for all AE-C and therefore the ROCL indifference prediction is supported. If we can reject the statistical null hypothesis of the generalized Fisher Exact test, we can conclude that at least in two of the AE-C pairs the proportions of subjects that chose the C lottery are different. For example, suppose there were only 2 AE-C lottery pairs. If the Fisher Exact test rejects the null hypothesis, we conclude that the proportions of subjects choosing the C lottery are not the same in the two lottery pairs. Therefore, even if one of the proportions was equal to 50%, as ROCL predicts for any given AE-C lottery pair, the rejection of the statistical null would imply that the other proportion is not statistically equal to 50%. Consequently, we would reject the research hypothesis that subjects satisfy ROCL and choose at random between the AE and the C lottery in all of the AE-C lottery pairs.

The generalized Fisher Exact test is appropriate to test the joint hypothesis that the proportion of subjects that chose the C lottery is the same in all of the AE-C lottery pairs. The basic assumptions of the test are satisfied by the data. As described by Sheskin[2004; p. 424 and 506], the assumptions are: (i) each of the *n* observations are independent (i.e., the outcome of one observation is not affected by the outcome of another observation), (ii) each observation is randomly selected from a population, (iii) each observation can be classified only into mutually exclusive categories, (iv) the Fisher Exact test is recommended when the size of the sample is small, and (v) many sources note that an additional assumption is that the sum of the rows and columns of the contingency table used in the Fisher Exact test are predetermined by the researcher.

Assumptions (i)-(iii) are satisfied for the same reasons we explained in the Binomial probability test. The Fisher Exact test is commonly used for small samples like ours instead of the Chi-square test of homogeneity, which relies on large samples to work appropriately. Finally, the last assumption is not met by our data, however, Sheskin [2004; p. 506] claim that this is rarely met in practice and, consequently, the test is used in contingency tables when "one or neither of the marginal sums is determined by the researcher."

1-in-40 Treatment

The strategy to test the basic prediction of indifference in the "Pay-1-in-40 compound is different from the one used in the "Pay-1-in-1-compound." The reason is that the structure of the data in each of the treatments is different. In the case of the 1-in-1 treatment, each of the 10 AE-C lottery pairs generated, using the terminology of the statistical literature, an "independent" sample in the sense that there was no subject that made choices over more than one AE-C lottery pair. On the contrary, in the 1-in-40 treatment we have multiple "dependent" samples. This means that several subjects made choices over each of our 10 AE-C lottery pairs, and therefore, we obtain 10 "dependent" samples. This subtle difference has relevant implications for the type of nonparametric test that one should use to test any hypothesis with the structure of the data we described. We use the Cochran Q test to test the basic ROCL prediction of indifference in the 1-in-40 treatment.

Statistical Null Hypothesis of the Cochran Q Test: The proportion of subjects that choose the C is the same in each of the 10 AE-C lottery pairs.

The information needed to perform this test is captured in a table of the following type:

	AE-C pair 1	AE-C pair 2		AE-C pair 9	AE-C pair 10
Subject 1	C ₁₋₁	c ₁₋₂	•	C ₁₋₉	C ₁₋₁₀
Subject 2	c ₂₋₁	$\mathcal{C}_{2\text{-}2}$		c_{2-9}	C ₂₋₁₀
	•				
Subject n	C _{n-1}	c_{n-2}		C _{n-9}	C_{n-10}
Total	$\sum_{i-1,2,\dots,n} \left(c_{i-1} \right)$	$\sum_{i-1,2,\dots,n} \left(c_{i-2} \right)$		$\sum_{i-1,2,\dots,n} \left(c_{i-9} \right)$	$\sum_{i-1,2,\dots,r} \left(c_{i-10} \right)$

This joint hypothesis is enough to reject the indifference prediction. For example, suppose there were only 2 AE-C lottery pairs. If the Cochran Q test rejects the null hypothesis, we conclude that the proportions of subjects choosing the C lottery are not the same in the two lottery pairs. Therefore, even if one of the proportions is equal to 50% as ROCL predicted, the test provides evidence that the other proportion is not equal to 50%. Consequently, we would reject the research hypothesis that subjects satisfy ROCL and choose at random between the AE and the C lottery in any of the AE-C lottery pairs. In the text we provide confidence intervals³⁶ on the number of subjects (out of the 62 in our sample) that chose the S lottery in each of the AE-C lottery. If for a given AE-C lottery pair the number 31 is not contained in the confidence interval, it implies that with 95% probability the proportion of the 62 subjects that chose the C lottery will not be 50%. We could have apply the Binomial probability test to each of the 10 AE-C pairs in the 1-in-40 treatment. However, this would not be appropriate since the Binomial test assumes "independence" in the sample in the statistical sense which is not satisfied in the present treatment.

The Cochran Q test is an appropriate test in this treatment because it allows us to jointly reject the null hypothesis that subjects choose the C lottery and the AE lottery in equal proportions when the data set is composed by multiple dependent samples. The basic assumptions of the test are satisfied by the data. As described by Sheskin[2004; p. 245], the assumptions are: (i) each of the subjects respond to each of the 10 AE-C lottery pairs, (ii) one has to control for order effects and (iii) each observation can be classified only into mutually exclusive categories. Assumptions (i) is

³⁶ We use the -total- Stata command to calculate the confidence intervals.

satisfied since in the 1-in-40 treatment each subjects respond to all 40 lottery pairs, which include the 10 AE-C pairs. Assumption (ii) is also satisfied because in our experiments each subject is presented with the 40 lotteries in random order. Assumption (iii) is trivially satisfied since in each of the AE-C pairs the subjects have to make a dichotomous choice.

B. Choice Patterns Where ROCL Predicts Consistent Choices

Research Null Hypothesis: Subjects choose the S lottery when presented with the S-C lottery pair if and only if they also choose the S lottery when presented with the corresponding S-AE lottery pair. This is equivalent to state the null hypothesis using the C and AE lotteries but we chose to work with the S lottery for simplicity. Therefore, ROCL is satisfied if we can provide statistical evidence that the proportion of subjects that choose the S lottery when presented with a S-C pair is equal to the proportion of subjects that choose also the S lottery when presented with the paired S-AE pair.

Structure of data sets: We use data from the 62 subjects in the 1-in-40 treatment who were presented with each of the 30 lottery pairs in Tables B2 and B3. Each of the 15 S-C lottery pairs in Table B2 has a corresponding S-AE pair in Table B3. Therefore, we can construct 15 comparisons of pairs that constitute 15 consistency tests of ROCL. In the 1-in-40 we again have to assume that the independence axiom holds. Therefore, we also use data from the 1-in-1 treatment to control for possible confounds created by the RLIM. However, we have to assume homogeneity in risk preferences for the analysis of this particular treatment. The reason is that the response of any subject to a particular S-C lottery pair, is going to be compared with the response of another subject to the paired S-AE lottery pair. In terms of the statistical literature, the responses to each of the S-C or S-AE pairs in the 1-in-1 constitute an "independent" sample. Conversely, the responses to each of the S-C subjects responded to each of the 15 S-C and the 15 S-AE. Also, each of the 15 comparisons is constructed by matching a S-C pair with its corresponding S-AE pair.

Analysis of data from the 1-in-1 treatment

We use the Fisher Exact test to evaluate the consistency predicted by ROCL in each of the paired comparisons of S-C pairs and S-AE pairs for which we have enough data to conduct the test. We also use the Cochran-Mantel-Haenszel (CMH) as a joint test of the 15 paired comparisons to evaluate the overall performance of the ROCL consistency prediction.

Statistical Null Hypothesis of the Fisher Exact Test: For any given paired comparison, subjects choose the S lottery in the same proportion when presented with a S-C pair and with its corresponding S-AE lottery pair.

The table shows statistical tests on contingency tables of the following form:

	Ch		
	S	AE/C	Total
S-AE pair	а	b	a+b
S-C pair	С	d	c+d
Total	a+c	b+d	a+b+c+d

The positions in each cell are defined as follows. The letter a represents the number of individuals that chose the simple lottery when they were presented with a S-AE pair. The letter c represents the number of individuals that chose the simple lottery when they were presented with the corresponding S-C pair. The letter b represents the number of subjects that chose the AE lottery when they were presented with the S-AE lottery pair. Similarly, the letter d represents the number of subjects that chose the C lottery when they were presented with the corresponding S-C lottery pair. In the notation of the previous note, the proportions used in the Fisher Exact test are defined as $\pi_1 = a/(a+b)$ and $\pi_2 = c/(c+d)$.

The Fisher Exact test for 2×2 contingency tables is appropriate to test individually in each of the 15 matched pairs the hypothesis that the proportion of subjects that chose the S lottery is the same when they are presented with the S-C or its corresponding S-AE pair. The basic assumptions of the test are satisfied by the data. As described by Sheskin[2004; p. 424 and 506], the assumptions are: (i) each of the n observations are independent (i.e., the outcome of one observation is not affected by the outcome of another observation), (ii) each observation is randomly selected from a population, (iii) each observation can be classified only into mutually exclusive categories, (iv) the Fisher Exact test is recommended when the size of the sample is small, and (v) many sources note that an additional assumption is that the sum of the rows and columns of the contingency table used in the Fisher Exact test are predetermined by the researcher. Assumptions (i)-(iv) are satisfied for the same reasons we explained in the in the case of the generalized Fisher exact test. Finally, as we explained before assumption (v) is not satisfied in our data.

Statistical Null Hypothesis of the Cochran-Mantel-Haenszel test: In all of the 15 paired comparisons subjects choose in the same proportion the S lottery when presented with the S-C lottery pair and its paired S-AE lottery pairs. More formally, the odds ratio of each of the 15 contingency tables constructed from the 15 paired comparisons are jointly equal to 1.

If the CMH test rejects the null hypothesis, then we interpret this as evidence of ROCL-inconsistent observed behavior. However, if we cannot reject the null, we conclude that subjects make choices according to the ROCL consistency predictions in the 15 pair comparisons even if we find that the Fisher Exact tests rejects its null hypothesis for some of the paired comparisons.

The CMH is the appropriate joint test to apply since it allows us to pool the data of multiple contingency tables that satisfy the assumptions needed for the Fisher Exact test and test jointly the homogeneity of the tables.

Analysis of data from the 1-in-40 treatment

We use the Cochran Q test coupled with the Bonferroni-Dunn (B-D) method to test the statistical hypothesis that subjects choose the S lottery in the same proportion when presented with a S-C lottery pair and with the corresponding S-AE lottery pair. The B-D method allows us to test if, in each of the 15 paired comparisons of S-C and S-AE lottery pairs, the observed difference in the proportion of subjects that chose the S lottery is statistically significant.

The B-D method is a *post-hoc* procedure that is conducted after calculating the Cochran Q test. The first step is to conduct the latter test to reject or not reject the null hypothesis that the proportions of individuals that choose the S lottery are the same in all 15 S-C and 15 S-AE lottery pairs. If this null is rejected the B-D method involves calculating a critical value d (see Sheskin[2004; p. 871] for the definition) that allows to evaluate the statistical significance of the difference in proportions and that takes into account all the information of the 30 lottery pairs and a confidence level of α .

Statistical Null Hypothesis in each of the Pair Comparisons using the B-D method. Define p_1 as the proportion of subjects that choose the S lottery when presented with a given S-AE lottery pair. Similarly, define p_2 as the proportion of subjects that chose the S lottery in the paired S-C lottery pair. The statistical null hypothesis is that, for a given paired comparison, $p_1 = p_2$.

The B-D method rejects the statistical null hypothesis if $|p_1-p_2| > d$. In this case we would conclude that the observed difference in proportions in a given paired comparison is statistically significant. This is a more powerful test than conducting individual tests for each paired comparison because the critical value d takes into account the information of all of the 15 comparisons. See Sheskin [2004; p. 871] for further details of the B-D method.

The Cochran Q test coupled with the B-D method are appropriate to test in this treatment the null hypothesis that subjects choose the S lottery in the same proportion when presented with a given S-C pair and with its corresponding S-AE lottery, and the data set is composed by multiple dependent samples in the sense we explained above. The basic assumptions of the Cochran Q test are satisfied by the data. As described by Sheskin [2004; p. 245], the assumptions are: (i) each of the subjects respond to each of the 15 S-C lottery pairs and the 15 S-AE lottery pairs, (ii) one has to control for order effects and (iii) each observation can be classified only into mutually exclusive categories. Assumptions (i) is satisfied since in the 1-in-40 treatment each subjects respond to all 40 lottery pairs, which include the 15 S-C pairs and the 15 S-AE pairs. Assumption (ii) is also satisfied because in our experiments each subject is presented with the 40 lotteries in random order. Assumption (iii) is trivially satisfied since in each of the AE-C pairs the subjects have to make a dichotomous choice. The B-D method applied to the Cochran Q test does not require any extra assumptions. However, the calculation of the critical value to make the comparisons requires to define a family wise Type I error rate (α_{FW}). Sheskin [2004; p. 871] claims that "[w]hen a limited number of comparisons are planned prior to collecting the data, most sources take the position that a researcher is not obliged to control the value of α_{FW} . In such a case, the per comparison Type I error rate (α_{PC}) will be equal to the prespecified value of alpha [the confidence level]."

We are also interested in studying the patterns in the violation of ROCL. We want to test the statistical validity of differences in switching behavior. A pattern inconsistent with ROCL would be

subjects choosing the S lottery when presented with a given S-C lottery pair, but switching to prefer the AE lottery when presented with the matched S-AE pair. We construct 2 × 2 contingency tables that show the number of subjects in any given matched pair who exhibit each of the four possible choice patterns: (i) always choosing the S lottery; (ii) choosing the S lottery when presented with a S-C pair and switching to prefer the AE lottery when presented with the matched S-AE pair; (iii) choosing the C lottery when presented with a S-C pair and switching to prefer the S lottery when presented with the matched S-AE; and (iv) choosing the C lottery when presented with the S-C lottery and preferring the AE lottery when presented with the matched S-AE. We use the McNemar test to evaluate the statistical significance of patterns in the violations of ROCL.

Statistical Null Hypothesis of the McNemar Test: Subjects exhibit the discordant choice patterns (ii) and (iii) in equal proportions within each set of matched pairs.

If the statistical null hypothesis is rejected then we can claim that there is an statistical difference in the two possible patterns of switching behavior that violate ROCL.

The test requires to construct a contingency tables of the following form:

	Simple Lottery vs. Actuarially- Equivalent			
		Left Lottery	Right Lottery	Total
	Left lottery	а	b	a+b
Simple Lottery vs.	Right lottery	C	d	c+d
Compound Lottery	Total	a+c	b+d	a+b+c+d

The positions in each cell are defined as follows. The letter *a* represents the number of individuals that chose the left lottery both when they were presented with a pair of a simple lottery and a compound lottery (S-C) *and* a corresponding pair that has the same simple lottery and the actuarially-equivalent lottery (S-AE) of the compound lottery. The simple lotteries, and therefore the compound and their actuarially equivalent lotteries, are always in the same position across. For the purpose of the statistical tests, the simple lotteries are always the left lotteries; the compound and their actuarially-equivalent lotteries are always the right lotteries. Therefore, *a* is the number of individuals that chose the simple lottery when the were presented with a given pair of S-C and its corresponding pair of S-AE. The letter *c* represents the number of individuals that chose the simple lottery when they were presented with a given pair of S-AE but that chose the compound lottery when they were presented with corresponding pair of S-C.

The McNemar test is an appropriate test to apply in this context. The assumptions of the test are (See Sheskin [2004; pp. 634]): (i) the sample of n subjects has been randomly selected from the population, (ii) each of the n onservations in the contingency table is independent of other

observations, (iii) each observation can be classified only into mutually exclusive categories, and (iv) the test should not be used with extremely small samples. Assumptions (i) and (iii) are satisfied for reasons we explained before. Even though there is no agreement of what an small sample is for the McNemar test, we follow the recommendation of the literature and provide in our results the exact probability of the test. Assumption (ii) is not satisfied since each subject respond makes choice over more than one pair. However, the test still allows us to draw conclusions about the discordant switching patterns but does not allow to make causal inferences, which is enough for our purposes. In fact it will allows us to conclude if there is an statistical difference between the two possible choice patterns that contradict ROCL Nevertheless, it will not allow us to conclude anything about the source of this difference (see Sheskin [2004; p. 639]).

Appendix E: Additional Econometric Analysis (NOT FOR PUBLICATION)

Analysis of Data from the 1-in-1 Treatment.

Assuming homogeneity in preferences, we find that the model that best describes the data is the one that allows for the source-dependent version of RDU, and conditional on this model there is no evidence of violations of ROCL. Both the Vuong test and the Clarke test provide statistical evidence that the model that allows for the source-dependent version of the RDU is the best model to explain the data in the 1-in-1 treatment.³⁷ Table E1 shows the parameter estimates of the two models we consider. In particular, panel A shows the estimates for the model that allows for the source-dependent version of RDU. We find that the estimates for parameters r, rc, y and yc are 0.62, 0.14, 0.77 and -0.19, respectively. A test of the joint null hypothesis that $rc = \gamma c = 0$ results in a pvalue of .29. This implies that there is no statistical evidence for source-dependency both in the utility and the probability weighting functions, and therefore no evidence of violations of ROCL when homogeneity is assumed. If we had assumed that subjects behave according to the sourcedependent EUT, we would have incorrectly concluded that there is evidence of violations of ROCL as suggested by the estimate of rc equal to 0.27 which has a p-value of 0.008 (see panel B of Table E1). A joint test of r and rc results in a p-value less than 0.0001. This highlights the importance of choosing the preference representation that best characterizes the way in which individuals make choices.

Although the model that best characterizes behavior in the 1-in-1 treatment is the source-dependent version of the RDU model, there is evidence of marginal diminishing returns but no evidence of probability weighting.³⁸ The parameter estimates for r and γ are equal to 0.62 and 0.77. Figure E1 shows the functions implied by these estimates and are plotted in the relevant domains. A test for the hypothesis that $\gamma=1$ results in a *p*-value of 0.23, which provides no evidence of probability weighting.

Analysis of Data from the 1-in-40 Treatment.

Assuming homogeneity, we find that the model that best describes the data is the one that allows for the source-dependence version of RDU, and conditional on this model we find evidence of violations of ROCL. Both the Vuong test and the Clarke test provide evidence to support the source-dependent RDU as the best model to explain the data in the 1-in-40 treatment.³⁹ An statistical test for the joint null hypothesis that $rc = \gamma c = 0$ results in a *p*-value less than 0.0001.

The estimates for r, rc, γ and γ c are 0.57, 0.11, 1.09 and -0.40, respectively. This implies that the nature of the violation has two components. First, the estimates suggests that when a typical

³⁷ When we assume homogeneity in risk attitudes, the Vuong test statistic is -1.45 in favor of the source-dependent RDU, with a *p*-value of .073. Further, the Clarke test also gives evidence in favor of the source-dependent RDU with a test statistic equal to 44.

³⁸ The Vuong and the Clarke tests provide evidence to choose the model that bests characterize data between two models but are agnostic about the statistical significance of the winning model.

³⁹ When we assume homogeneity in risk attitudes, the Vuong test statistic is -5.16 in favor of the source-dependent RDU, with a *p*-value less than 0.001. Further, the Clarke test also gives evidence in favor of the source-dependent RDU with a test statistic equal to 935.

individual is presented with a compound lottery he behaves as if the utility function was more concave. The linear combination of parameters r + rc results in an estimated coefficient of 0.68 with a *p*-value less than 0.0001 and a test of the null hypothesis of r + rc = 0.57 results in a *p*-value equal to 0.0001. Thus a typical subject would increase his utility risk aversion parameter from 0.57 to 0.68 when presented with a compound lottery. Second, there is no evidence of probability weighting when subjects are presented with simple lotteries but there is evidence of probability optimism when subjects evaluate compound lotteries. A test on the probability weighting parameter for $\gamma = 1$ results in a *p*-value equal to 0.28 and the linear combination $\gamma + \gamma c$ results in an estimated parameter equal to 0.69 with a *p*-value less than 0.001. Hence, when presented with a simple lottery a typical subject displays no probability weighting but does exhibits diminishing marginal returns; however, when facing a compound lottery, a typical subject behaves as if the utility function was more concave and the probability weighting function displays probability optimism. Figure E2 shows how the concavity of the utility function and the probability weighting function differ when individuals are presented with a simple or a compound lottery.

Table E1: Estimates of Source-Dependent RDU and EUT Model Allowing for Heterogeneity

Data from the 1-in-1 treatment (N=133). Estimates from the Fechner error parameter omitted

A. Source-Dependent RDU(LL=-81.82)

	Coef.	Std. Err.	Z	P> z	[95% Conf.	Interval]
r	. 6182909	. 0669022	9. 24	0.000	. 4871649	. 7494169
r_rocl	. 1410226	. 0922965	1. 53	0. 127	0398752	. 3219204
γ	. 7699848	. 193429	3. 98	0.000	. 3908708	1. 149099
γC	1955013	. 1819329	-1.07	0. 283	5520832	. 1610806


(Ho: $rc = \gamma c = 0$; p-value = 0.2848) (Ho: $r = \gamma = 0$; p-value < 0.0001)

B. Source-Dependent Version of EUT (LL=-84.24)

	Coef.	Robust Std. Err.	z	P> z	[95% Conf.	Interval]
r	. 5799532	. 0731694	7. 93	0.000	. 4365438	. 7233626
rc	. 2717936	. 1028005	2.64	0.008	. 0703083	. 4732789

(Ho: r=rc=0; p-value < 0.0001)

Figure E1: Estimated Functions from the RDU Specification in the 1-in-1 Treatment Assuming Homogeneity in Preferences






Table E2: Estimates of Source-Dependent RDU and EUT Model Allowing for Heterogeneity

Data from the 1-in-40 treatment (N=2480=62 Subjects \times 40 choices). Estimates from the Fechner error parameter omitted

A. Source-Dependent RDU (LL=-1460.57)


	Coef.	Robust Std. Err.	Z	P> z	[95% Conf.	Interval]
r	. 5670508	. 024286	23. 35	0. 000	. 519451	. 6146505
rc	. 1090157	. 0231279	4. 71	0.000	. 0636858	. 1543456
γ	1. 088315	. 081601	13. 34	0. 000	. 9283802	1. 24825
γС	 3957716	. 0587485	-6. 74	0.000	5109165	2806266

(Ho: $rc = \gamma c = 0$; p-value<0.0001) (Ho: $r = \gamma = 0$; p-value=0.0001)

B. Version of EUT (LL=-1512.45)

	Coef.	Std. Err.	z	P> z	[95% Conf.	Interval]
r	. 5906296	. 0352857	16. 74	0. 000	. 5214708	. 6597883
rc	0071859	. 0194071	-0. 37	0. 711	0452233	. 0308514

Figure E2: Estimated Functions from the RDU Specification in the 1-in-40 Treatment Assuming Homogeneity in Preferences



Appendix F: Detailed Binomial Test Results (NOT FOR PUBLICATION)

The following tests are conducted over the actuarially-equivalent and compound lottery pairs in Table B4 (Pairs 31-40).

<u>Binomial tests for treatment 1-in-1</u>: We do not report the test for pairs 31, 35 and 35 because for the first one there was only one observation and for the last two there were no observations at all.

Note: In the following Binomial tests "N" is the total size of the sample for each lottery pair and "Observed k" is the number of subjects (out of the N subjects) that chose the compound lottery.

Pair 32

Pair 33

Variable	N	Observed k	Expected k	Assumed p	Observed p
choice_aed~3	7	5	3.5	0.50000	0.71429
Pr(k >= 5) Pr(k <= 5) Pr(k <= 2 or k >=	=	0.937500 (one-sided test) one-sided test)		

Pair 36

Pair 37

Pair 38

<u>Pair 39</u>

Variable	N	Observed k	Expected k	Assumed p	Observed p
choice_aed~9	5	4	2.5	0.50000	0.80000
Pr(k >= 4) Pr(k <= 4) Pr(k <= 1 or k >	=	0.968750 (o	ne-sided test) ne-sided test) wo-sided test)		

Pair 40

Variable	N	Observed k	Expected k	Assumed p	Observed p
choice_ae~10	6	3	3	0.50000	0.50000
Pr(k >= 3) Pr(k <= 3) Pr(k <= 3 or k >	=	0.656250 (o	ne-sided test) ne-sided test) wo-sided test)		

Appendix G: Detailed Fisher Exact Test Results (NOT FOR PUBLICATION)

The following tests are conducted by matching each of the 15 simple and compound lottery pairs in Table B2 (Pairs 1-15) with a corresponding pair in the set of the 15 simple and actuarially-equivalent lottery pairs in Table B3 (Pairs 16-30).

Pair 1 vs. Pair 16

		Choice	
Pairs	S	C/AE	Total
S-AE Pair 16 S-C Pair 1	3 1	0 1	3 2
Total	4	1	5
	sher's exact		0.400 0.400

Pair 3 vs. Pair 18

	Choice	e	
Pairs	S	C/AE	Total
S-AE Pair 18 S-C Pair 3	3 0	3 2	6 2
Total	3	5	8
	isher's exact = isher's exact =		0.464 0.357

Pair 5 vs. Pair 20

	Cho	oice	
Pairs	S	C/AE	Total
S-AE Pair 18 S-C Pair 5	0 2	1 0	1 2
Total	2	1	3
	isher's exact = isher's exact =		0.333 0.333

Pair 6 vs. Pair 21

	Choice				
Pairs	S	C/AE	Total		
S-AE Pair 21 S-C Pair 6	2 2 2	1 2	+3 3 4		
Total	4	3	7		

Fisher's exact = 1.000 1-sided Fisher's exact = 0.629

Pair 7 vs. Pair 22

	Choice			
Pairs	S	C/AE	Total	
S-AE Pair 22 S-C Pair 7	4 5	0 4	4 9	
Total	9	4	13	

Fisher's exact = 0.228 1-sided Fisher's exact = 0.176

Pair 8 vs. Pair 23

	Choice			
Pairs	S	C/AE	Total	
S-AE Pair 23 S-C Pair 8	1 2	2 2	3 4	
Total	3	4	7	

Fisher's exact = 1.000 1-sided Fisher's exact = 0.629

Pair 9 vs. Pair 24

		Choice	
Pairs	S	C/AE	Total
S-AE Pair 24 S-C Pair 9	0 5	3 1	3 6
Total	5	4	9

Fisher's exact = 0.048 1-sided Fisher's exact = 0.048

Pair 11 vs. Pair 26

	Choice		
Pairs	S	C/AE	Total
S-AE Pair 26 S-C Pair 11	3 5	2 4	+ 5 9
Total	8	6	14

Fisher's exact = 1.000 1-sided Fisher's exact = 0.657

Pair 12 vs. Pair 27

	Choice		
Pairs	S	C/AE	Total
S-AE Pair 27 S-C Pair 11	4 2	1 0	+ 5 2
Total	6	1	7

Fisher's exact = 1.000 1-sided Fisher's exact = 0.714

Pair 13 vs. Pair 28

	Choice		
Pairs	S	C/AE	Total
S-AE Pair 28 S-C Pair 13	2	2 0	4 1
Total	3	2	5

Fisher's exact = 1.000 1-sided Fisher's exact = 0.600

Pair 15 vs. Pair 30

	Choice		
Pairs	S	C/AE	Total
S-AE Pair 30 S-C Pair 15	3	0 1	3 1
Total	3	1	4

Fisher's exact = 0.250
1-sided Fisher's exact = 0.250

Appendix H: Detailed McNemar Test Results (NOT FOR PUBLICATION)

All Simple-Compound pairs vs. all Simple-Actuarially-Equivalent pairs

Cases	Controls Exposed	Unexposed	Total
Exposed Unexposed	447 182	97 204	544 386
Total	 629	301	930

McNemar's chi2(1) = 25.90 Prob > chi2 = 0.0000 Exact McNemar significance probability = 0.0000

Proportion with factor

Cases	.5849462			
Controls	.6763441	[95% Conf.	<pre>Interval]</pre>	
difference	0913978	1271815	0556142	
ratio	.8648649	.8177913	.9146481	
rel. diff.	282392	405559	1592251	
odds ratio	.532967	.4121939	.6855842	(exact)

Note: the contingency table above was constructed by summing each of the positions in the 15 contingency tables below. For example, the number in the first position (column 1, row 1) in the table above was calculated by summing the numbers in the first position in each of the 15 contingency tables below.

Pair 1 vs. Pair 16

Cases	Controls Exposed	Unexposed	 Total
Exposed Unexposed	22 32	2 6	24 38
Total	54	8	

McNemar's chi2(1) = 26.47 Prob > chi2 = 0.0000 Exact McNemar significance probability = 0.0000

Cases	.3870968		
Controls	.8709677	[95% Conf. Interval]	
difference	483871	63953853282034	
ratio	.444444	.3235546 .6105024	
rel. diff.	-3.75	-6.8634686365323	
odds ratio	.0625	.0072572 .2449784 (e:	xact)

Pair 2 vs. Pair 17

Cases	Controls Exposed	Unexposed	Total
Exposed Unexposed	58	1	59 3
Total	61	1	62

McNemar's chi2(1) = 1.00 Prob > chi2 = 0.3173 Exact McNemar significance probability = 0.6250

Proportion with factor

Cases	.9516129			
Controls	.983871	[95% Conf.	<pre>Interval]</pre>	
difference	0322581	1110998	.0465837	
ratio	.9672131	.9060347	1.032522	
rel. diff.	-2	-8.789514	4.789514	
odds ratio	.3333333	.0063495	4.151441	(exact)

Pair 3 vs. Pair 18

Cases	Controls Exposed	Unexposed	Total
Exposed	38	1	39
Unexposed	17	6	23
Total			
	55	7	62

McNemar's chi2(1) = 14.22 Prob > chi2 = 0.0002 Exact McNemar significance probability = 0.0001

Cases	.6290323			
Controls	.8870968	[95% Conf.	<pre>Interval]</pre>	
difference	2580645	3919297	1241993	
ratio	.7090909	.5925527	.8485489	
rel. diff.	-2.285714	-4.438998	1324308	
odds ratio	.0588235	.0014075	.3754091	(exact)

Pair 4 vs. Pair 19

Cases	Controls Exposed	Unexposed	Total
Exposed Unexposed	7 7	6 42	13
Total	14	48	62

McNemar's chi2(1) = 0.08 Prob > chi2 = 0.7815 Exact McNemar significance probability = 1.0000

Proportion with factor

Cases Controls	.2096774	[95% Conf.	Interval]	
difference ratio rel. diff.	.9285714	1461672 .5499486 169583	.1139091 1.567864 .1279163	
odds ratio	.8571429	.2379799	2.978588	(exact)

Pair 5 vs. Pair 20

Cases	Controls Exposed	Unexposed	 Total
Exposed Unexposed	12 13	6 31	18 44
Total	25	37	62

McNemar's chi2(1) = 2.58 Prob > chi2 = 0.1083 Exact McNemar significance probability = 0.1671

Cases	.2903226			
Controls	.4032258	[95% Conf.	<pre>Interval]</pre>	
difference	1129032	2639309	.0381244	
ratio	.72	.4813126	1.077055	
rel. diff.	1891892	4409851	.0626067	
odds ratio	.4615385	.1438515	1.301504	(exact)

Pair 6 vs. Pair 21

Cases	Controls Exposed	Unexposed	Total
Exposed Unexposed	35 11	6 10	41 21
Total	46	16	62

McNemar's chi2(1) = 1.47 Prob > chi2 = 0.2253 Exact McNemar significance probability = 0.3323

Proportion with factor

Cases	.6612903			
Controls	.7419355	[95% Conf.	<pre>Interval]</pre>	
difference	0806452	2255601	.0642697	
ratio	.8913043	.7399666	1.073594	
rel. diff.	3125	8911317	.2661317	
odds ratio	.5454545	.1656336	1.609034	(exact)

Pair 7 vs. Pair 22

Cases		Controls Exposed	Unexposed	 Total
Expos Unexpos		26 16	8 12	34 28
Tot	al	42	20	- 62

McNemar's chi2(1) = 2.67 Prob > chi2 = 0.1025 Exact McNemar significance probability = 0.1516

Cases Controls	.5483871 .6774194	[95% Conf.	Interval]	
difference ratio rel. diff.	1290323 .8095238 4	2966623 .6278837 9680515	.0385978 1.04371 .1680515	
odds ratio	.5	.1852586	1.238236	(exact)

Pair 8 vs. Pair 23

Cases	Controls Exposed	Unexposed	Total
Exposed Unexposed	23 11	11 17	34 28
Total	34	28	62

McNemar's chi2(1) = 0.00 Prob > chi2 = 1.0000 Exact McNemar significance probability = 1.0000

Proportion with factor

Cas Con	es trols	.5483871 .5483871	[95% (Conf.	Interval]	
rat	ference io . diff.	0 1 0	164 .7630 3283	0866	.164404 1.310467 .3283231	
odd	s ratio	1	.3933	1661	2.543454	(exact)

Pair 9 vs. Pair 24

Cases	Controls Exposed	Unexposed	 Total +
Exposed Unexposed	I	6 10	43 19
Total	46	16	62

McNemar's chi2(1) = 0.60 Prob > chi2 = 0.4386 Exact McNemar significance probability = 0.6072

Cases Controls	.6935484 .7419355	[95% Conf.	Interval]	
difference ratio rel. diff.	.9347826	1863563 .7881077 7045	.0895821 1.108755 .3295	
odds ratio	.6666667	.1952634	2.097224	(exact)

Pair 10 vs. Pair 25

Cases		Controls Exposed	Unexposed	Total
	Exposed Unexposed	45 12	3 2	48 14
	Total	 57	5	62

McNemar's chi2(1) = 5.40 Prob > chi2 = 0.0201 Exact McNemar significance probability = 0.0352

Proportion with factor

Cases Controls	.7741935 .9193548	[95% Conf.	Interval]	
difference ratio rel. diff.	1451613 .8421053 -1.8		0120515 .9736266 .7404037	
odds ratio	.25	.0452729	.9263782	(exact)

Pair 11 vs. Pair 26

Cases	 	Controls Exposed	Unexposed	 Total
Uı	Exposed nexposed	27 9	11 15	38 24
	Total	36	26	62

McNemar's chi2(1) = 0.20 Prob > chi2 = 0.6547 Exact McNemar significance probability = 0.8238

Cases	.6129032			
Controls	.5806452	[95% Conf.	<pre>Interval]</pre>	
difference	.0322581	1250174	.1895335	
ratio	1.055556	.832837	1.337834	
rel. diff.	.0769231	2469752	.4008214	
odds ratio	1.222222	.460447	3.336929	(exact)
odds ratio	1.222222	.460447	3.336929	(exact)

Pair 12 vs. Pair 27

Cases		Controls Exposed	Unexposed	Total
	Exposed Unexposed	26 9	14 13	40 22
	Total	35	27	62

McNemar's chi2(1) = 1.09 Prob > chi2 = 0.2971 Exact McNemar significance probability = 0.4049

Proportion with factor

Cases	.6451613	
Controls	.5645161	[95% Conf. Interval]
difference	.0806452	0857564 .2470467
ratio	1.142857	.888976 1.469244
rel. diff.	.1851852	1290666 .4994369
odds ratio	1.555556	.6271247 4.074174 (

odds ratio 1.555556 .6271247 4.074174 (exact)

Pair 13 vs. Pair 28

Cases		Controls Exposed	Unexposed	Total
	Exposed Unexposed	44	1 7	45 17
	Total	54	8	62

McNemar's chi2(1) = 7.36 Prob > chi2 = 0.0067 Exact McNemar significance probability = 0.0117

Cases Controls	.7258065 .8709677	[95% Conf.	Interval]	
difference	1451613	2597136	030609	
ratio	.8333333	.7303802	.9507986	
rel. diff.	-1.125	-2.309497	.0594969	
odds ratio	.1	.0023043	.702939	(exact)

Pair 14 vs. Pair 29

Cases	 +	Controls Exposed	Unexposed	 Total
Expos Unexpos		34 12	8 8	42 20
Tota	al	46	16	62

McNemar's chi2(1) = 0.80 Prob > chi2 = 0.3711 Exact McNemar significance probability = 0.5034

Proportion with factor

LOII WICII LAC	J C O I			
Cases	.6774194			
Controls	.7419355	[95% Conf.	<pre>Interval]</pre>	
difference	0645161	2211047	.0920725	
ratio	.9130435	.7479736	1.114542	
rel. diff.	25	8624887	.3624887	
odds ratio	.6666667	.2363844	1.773597	(exact)

Pair 15 vs. Pair 30

Cases		Controls Exposed	Unexposed	Total
	Exposed Unexposed	13 11	13 25	26 36
	Total	 24	38	62

McNemar's chi2(1) = 0.17 Prob > chi2 = 0.6831 Exact McNemar significance probability = 0.8388

Cases Controls	.4193548 .3870968	[95% Conf.	Interval]	
difference ratio rel. diff.	.0322581 1.083333 .0526316	1385308 .7376116 1933086	.2030469 1.591096 .2985718	
odds ratio	1.181818	.4885561	2.913432	(exact)

Appendix I: Detailed Wald Test Results for Predictions (NOT FOR PUBLICATION)

1-in-1 Treatment: Wald tests for the null hypothesis of r = 0

Predicted r

	Predicted r																
<i>p</i> -value	0.45	0.48	0.49	0.53	0.59	0.62	0.64	0.67	0.75	0.78	0.80	0.83	0.89	0.93	0.94	0.97	# of Subjects
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	28	0	28
1.14E-34	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	16	16
1.33E-26	0	0	0	0	0	0	0	0	0	0	0	0	22	0	0	0	22
5.60E-25	0	0	0	0	0	0	0	9	0	0	0	0	0	0	0	0	9
6.78E-25	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	2
2.90E-22	0	0	0	0	0	0	17	0	0	0	0	0	0	0	0	0	17
2.02E-19	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	9
7.87E-19	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	2
1.12E-15	0	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0	8
2.89E-13	0	0	0	0	0	5	0	0	0	0	0	0	0	0	0	0	5
3.05E-11	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	2
3.12E-09	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	3
4.42E-07	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	3
6.11E-07	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	2
0.0002934	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4
0.0003798	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
# of Subjects	4	1	2	3	8	5	17	9	2	3	2	2	22	9	28	16	133

1-in-1 Treatment: Wald tests for the null hypothesis of gamma = 1

Predicted gamma

									0								
<i>p</i> -value	0.23	0.24	0.25	0.26	0.29	0.30	0.31	0.32	0.46	0.47	0.48	0.49	0.52	0.53	0.54	0.55	# of Subjects
7.02E-06	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	22
0.000047	0	0	0	0	0	0	0	0	0	17	0	0	0	0	0	0	17
0.0000525	0	28	0	0	0	0	0	0	0	0	0	0	0	0	0	0	28
0.0001667	0	0	0	0	0	0	0	0	8	0	0	0	0	0	0	0	8
0.0002153	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	2
0.0004045	0	0	0	0	0	0	0	0	0	0	0	9	0	0	0	0	9
0.000434	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	2
0.0004769	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	2
0.0005309	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	3
0.0007535	0	0	9	0	0	0	0	0	0	0	0	0	0	0	0	0	9
0.001047	0	0	0	16	0	0	0	0	0	0	0	0	0	0	0	0	16
0.0022308	0	0	0	0	0	0	0	0	0	0	5	0	0	0	0	0	5
0.0147585	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	3
0.0306056	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
0.0315525	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	2
0.0396612	0	0	0	0	0	0	0	0	0	0	0	0	4	0	0	0	4
# of Subjects	22	28	9	16	2	2	3	2	8	17	5	9	4	2	1	3	133

1-in-40 Treatment: Wald tests for the null hypothesis of r = 0

Predicted r

<i>p</i> -value	0.40	0.44	0.47	0.49	0.52	0.62	0.64	0.67	0.67	0.71	0.82	0.86	# of Subjects
1.96E-27	0	0	0	0	0	0	7	0	0	0	0	0	7
1.6E-24	0	0	0	0	0	0	0	0	0	0	8	0	8
1.56E-20	0	0	0	0	0	0	0	0	3	0	0	0	3
2.35E-18	0	0	0	0	0	0	0	0	0	11	0	0	11
1.25E-15	0	0	0	0	8	0	0	0	0	0	0	0	8
5.45E-14	0	0	0	0	0	0	0	4	0	0	0	0	4
3.92E-13	0	0	0	0	0	0	0	0	0	0	0	3	3
2.07E-10	0	0	0	11	0	0	0	0	0	0	0	0	11
8.76E-10	0	0	0	0	0	1	0	0	0	0	0	0	1
1.22E-08	0	0	3	0	0	0	0	0	0	0	0	0	3
0.000000917	0	1	0	0	0	0	0	0	0	0	0	0	1
0.0000284	2	0	0	0	0	0	0	0	0	0	0	0	2
# of Subject	2	1	3	11	8	1	7	4	3	11	8	3	62

1-in-40 Treatment: Wald tests for the null hypothesis of r + rc = 0

						Predict	r + rc						
p-value	0.54	0.57	0.59	0.63	0.66	0.67	0.71	0.73	0.74	0.76	0.81	0.84	# of Subjects
0.00E+00	0	0	0	0	0	0	0	3	0	0	0	0	3
9.27E-32	0	0	0	0	0	0	7	0	0	0	0	0	7
2.86E-25	0	0	0	0	0	0	0	0	4	0	0	0	4
3.55E-22	0	0	0	0	0	0	0	0	0	11	0	0	11
5.59E-19	0	0	0	0	0	0	0	0	0	0	0	3	3
7.64E-18	0	0	0	0	0	0	0	0	0	0	8	0	8
2.03E-17	0	0	0	11	0	0	0	0	0	0	0	0	11
7.91E-14	0	1	0	0	0	0	0	0	0	0	0	0	1
2.32E-12	0	0	0	0	0	1	0	0	0	0	0	0	1
1.5E-10	0	0	0	0	8	0	0	0	0	0	0	0	8
0.000000094	0	0	3	0	0	0	0	0	0	0	0	0	3
0.000000886	2	0	0	0	0	0	0	0	0	0	0	0	2
# of Subject	2	1	3	11	8	1	7	3	4	11	8	3	62

1-in-40 Treatment: Wald tests for the null hypothesis of $\gamma = 1$

Predicted gamma

						1 icaici	ca gairiiri	ı					
<i>p</i> -value	0.37	0.41	0.68	0.75	0.79	0.95	0.99	1.06	1.26	1.30	1.33	1.37	# of Subjects
7.41E-03	0	8	0	0	0	0	0	0	0	0	0	0	8
0.0110372	3	0	0	0	0	0	0	0	0	0	0	0	3
0.0924481	0	0	0	0	0	0	0	0	0	0	0	11	11
0.1921943	0	0	0	0	0	0	0	0	0	0	8	0	8
0.1973165	0	0	0	11	0	0	0	0	0	0	0	0	11
0.2324818	0	0	1	0	0	0	0	0	0	0	0	0	1
0.2682751	0	0	0	0	0	0	0	0	0	2	0	0	2
0.3140454	0	0	0	0	0	0	0	0	1	0	0	0	1
0.3357557	0	0	0	0	3	0	0	0	0	0	0	0	3
0.8049443	0	0	0	0	0	0	0	3	0	0	0	0	3
0.8290983	0	0	0	0	0	4	0	0	0	0	0	0	4
0.9432108	0	0	0	0	0	0	7	0	0	0	0	0	7
# of Subjects	3	8	1	11	3	4	7	3	1	2	8	11	62

1-in-40 Treatment: Wald tests for the null hypothesis of $\gamma + \gamma c = 1$

Predicted gamma + gammac

<i>p</i> -value	0.29	0.34	0.42	0.55	0.56	0.61	0.61	0.63	0.69	0.69	0.82	0.87	# of Subjects
4.02E-05	0	0	0	7	0	0	0	0	0	0	0	0	7
0.0001762	0	0	1	0	0	0	0	0	0	0	0	0	1
0.0002076	8	0	0	0	0	0	0	0	0	0	0	0	8
0.008463	0	3	0	0	0	0	0	0	0	0	0	0	3
0.0122949	0	0	0	0	3	0	0	0	0	0	0	0	3
0.0125595	0	0	0	0	0	0	11	0	0	0	0	0	11
0.0382169	0	0	0	0	0	4	0	0	0	0	0	0	4
0.062012	0	0	0	0	0	0	0	0	0	3	0	0	3
0.0686239	0	0	0	0	0	0	0	0	1	0	0	0	1
0.0960365	0	0	0	0	0	0	0	2	0	0	0	0	2
0.2874661	0	0	0	0	0	0	0	0	0	0	11	0	11
0.3762546	0	0	0	0	0	0	0	0	0	0	0	8	8
# of Subject	8	3	1	7	3	4	11	2	1	3	11	8	62